

Results of study

FUJII, Shinobu

We are interested in isoparametric hypersurfaces in spheres with four distinct principal curvatures. We expect that every isoparametric hypersurface in spheres with four distinct principal curvatures is related to moment map.

A hypersurface in the spheres S^n is called an *isoparametric hypersurface* if this one is a level set of an isoparametric function on S^n . It is known that isoparametric functions on S^n are obtained by the restrictions on S^n of Cartan-Münzner polynomial functions $\varphi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$.

Let G/K be an irreducible Hermitian symmetric space of compact type and of rank two. Then we get two invariant functions under the isotropy representation of G/K . One comes from a Cartan-Münzner polynomial. It is known that a principal orbit of the isotropy representation of G/K is a homogeneous isoparametric hypersurface in a sphere. By the definition of isoparametric hypersurfaces, there exists an isoparametric function on the sphere. This function is the restriction of a Cartan-Münzner polynomial φ . This polynomial φ is invariant under the isotropy representation of G/K . Another invariant function comes from a moment map. Since G/K is Hermitian, the isotropy representation of G/K is a Hamiltonian action. Thus, there exists a moment map μ for this action. By definition, μ is equivariant under the action. Hence, a composition function of μ and a norm which is invariant under the isotropy representation of G/K is invariant under the action.

In [1], we proved that there exists a K -invariant norm on \mathfrak{k}^* such that φ coincides with the squared-norm of the moment map μ for the isotropy representation of G/K in the case of compact irreducible Hermitian symmetric spaces of rank two and of classical type.

Cartan-Münzner polynomials which we got are essentially the same as ones which are computed in Ozeki-Takeuchi (1976). The difference of both is a computational method.

We also showed that there exists a K -invariant norm on \mathfrak{k}^* such that φ coincides with the squared-norm of the moment map μ for the isotropy representation of G/K in the case of compact irreducible Hermitian symmetric spaces of rank two. This is based on joint work with Hiroshi Tamaru (Hiroshima University).