

# Uniqueness of an orthogonal 2-handle pair on a surface-link

Akio Kawauchi

*Osaka Central Advanced Mathematical Institute, Osaka Metropolitan University*

*Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan*

*kawauchi@osaka-cu.ac.jp*

*Keywords:* Surface-link, Orthogonal 2-handle pair. *Mathematics Subject Classification 2010:* Primary 57Q45; Secondary 57N13

## ABSTRACT

The proof of uniqueness of an orthogonal 2-handle pair on a surface-link is given from the viewpoint of a normal form of 2-handle core disks. A version to an immersed orthogonal 2-handle pair on a surface-link is also observed.

## 1. Introduction

A *surface-link* is a closed oriented (possibly disconnected) surface  $F$  embedded in the 4-space  $\mathbf{R}^4$  by a smooth (or a piecewise-linear locally flat) embedding. When  $\mathbf{F}$  is connected, it is also called a *surface-knot*. Two surface-links  $F$  and  $F'$  are *equivalent* by an *equivalence*  $f$  if  $F$  is sent to  $F'$  orientation-preservingly by an orientation-preserving diffeomorphism (or piecewise-linear homeomorphism)  $f : \mathbf{R}^4 \rightarrow \mathbf{R}^4$ . A *trivial* surface-link is a surface-link  $F$  which is the boundary of disjoint handlebodies smoothly embedded in  $\mathbf{R}^4$ , where a handlebody is a 3-manifold which is a 3-ball, a solid torus or a boundary-disk sum of some number of solid tori. A trivial surface-knot is also called an *unknotted* surface-knot and a trivial disconnected surface-link is also called an *unknotted and unlinked* surface-link. A trivial surface-link is unique up to

equivalences (see [1]). A *2-handle* on a surface-link  $F$  in  $\mathbf{R}^4$  is an embedded 2-handle  $D \times I$  on  $F$  with  $D$  a core disk such that  $D \times I \cap F = \partial D \times I$ , where  $I$  denotes a closed interval containing 0 and  $D \times 0$  is identified with  $D$ . If  $D$  is an immersed disk, then call it an *immersed 2-handle*. Two (possibly immersed) 2-handles  $D \times I$  and  $E \times I$  on  $F$  are *equivalent* if there is an equivalence  $f : \mathbf{R}^4 \rightarrow \mathbf{R}^4$  from  $F$  to itself such that the restriction  $f|_F : F \rightarrow F$  is the identity map and  $f(D \times I) = E \times I$ . An *orthogonal 2-handle pair* (or simply, an *O2-handle pair*) on  $F$  is a pair  $(D \times I, D' \times I)$  of 2-handles  $D \times I, D' \times I$  on  $F$  such that

$$D \times I \cap D' \times I = \partial D \times I \cap \partial D' \times I$$

and  $\partial D \times I$  and  $\partial D' \times I$  *meet orthogonally* on  $F$ , that is, the boundary circles  $\partial D$  and  $\partial D'$  meet transversely at one point  $q$  and the intersection  $\partial D \times I \cap \partial D' \times I$  is homeomorphic to the square  $Q = q \times I \times I$  (see [2, Fig.1]). An important property of an O2-handle pair  $(D \times I, D' \times I)$  on a surface-link  $F$  is the following property (see [2] for the proof):

**Common 2-handle property** Let  $F$  be a surface-link in  $\mathbf{R}^4$ , and  $(D \times I, D' \times I)$  and  $(E \times I, E' \times I)$  O2-handle pairs on  $F$  in  $\mathbf{R}^4$  with  $\partial D \times I = \partial E \times I$  and  $\partial D' \times I = \partial E' \times I$ . If  $D \times I = E \times I$  or  $E' \times I = D' \times I$ , then the O2-handle pairs  $(D \times I, D' \times I)$  and  $(E \times I, E' \times I)$  on  $F$  are equivalent by an equivalence obtained by 3-cell moves on the unions  $D \times I \cup D' \times I$  and  $E \times I \cup E' \times I$  which are 3-balls.

In this paper, the following uniqueness theorem of an O2-handle pair on a surface-link is shown by using a normal form of 2-handle core disks discussed in [4] and Common 2-handle property stated above repeatedly which is announced in [2, Section 3] with incomplete proof although the tools of the present proof appear there.

**Theorem 1.1.** Let  $F$  be a surface-link in  $\mathbf{R}^4$ , and  $(D \times I, D' \times I)$  and  $(E \times I, E' \times I)$  O2-handle pairs on  $F$  in  $\mathbf{R}^4$  with  $\partial D \times I = \partial E \times I$  and  $\partial D' \times I = \partial E' \times I$ . Then the O2-handle pairs  $(D \times I, D' \times I)$  and  $(E \times I, E' \times I)$  on  $F$  are equivalent.

This theorem for a trivial surface-link is heavily used for confirming the smooth unknotting conjecture of a surface-knot in [2] and the smooth unknotting-unlinking conjecture for a surface-link in [3]. For an immersed O2-handle pair, the following proposition is provided:

**Proposition 1.2.** If  $(D \times I, D' \times I)$  is an immersed O2-pair on a surface-link  $F$  in  $\mathbf{R}^4$  with  $D \times I$  immersed and  $D' \times I$  embedded, then there is an embedded 2-handle  $D_* \times I$  with  $\partial D_* \times I = \partial D \times I$  such that  $(D_* \times I, D' \times I)$  is an O2-handle pair on  $F$ .

For the proof of Proposition 1.2, Finger move canceling operation is used to cancel a double point of an immersed core disk  $D$  of the immersed 2-handle  $D \times I$  on  $F$ , which is explained in Section 3. By Theorem 1.1 and Proposition 1.2, we have the following corollary.

**Corollary 1.3.** Let  $F$  be a surface-link in  $\mathbf{R}^4$ , and  $(D \times I, D' \times I)$  and  $(E \times I, E' \times I)$  immersed O2-handle pairs on  $F$  in  $\mathbf{R}^4$  with  $\partial D \times I = \partial E \times I$  and  $\partial D' \times I = \partial E' \times I$ .

(1) If  $D' \times I$  and  $E' \times I$  are embedded, then there are embedded 2-handles  $D_* \times I$  and  $E_* \times I$  on  $F$  with  $\partial D_* \times I = \partial D \times I$  and  $\partial E_* \times I = \partial E \times I$  such that  $(D_* \times I, D' \times I)$  and  $(E_* \times I, E' \times I)$  are equivalent O2-handle pairs on  $F$ , so that the surface-links  $F(D' \times I)$  and  $F(E' \times I)$  are equivalent.

(2) If  $D' \times I$  and  $E \times I$  are embedded, then there are embedded 2-handles  $D_* \times I$  and  $E'_* \times I$  on  $F$  with  $\partial D_* \times I = \partial D \times I$  and  $\partial E'_* \times I = \partial E' \times I$  such that  $(D_* \times I, D' \times I)$  and  $(E \times I, E'_* \times I)$  are equivalent O2-handle pairs on  $F$ , so that the surface-links  $F(D' \times I)$  and  $F(E \times I)$  are equivalent.

The proof of Theorem 1.1 is done in Section 2 and the proof of Proposition 1.2 is done in Section 3. Throughout the paper, the notation

$$XJ = \{(x, t) \in \mathbf{R}^4 \mid x \in X, t \in J\}$$

is used for a subspace  $X$  of  $\mathbf{R}^3$  and a subinterval  $J$  of  $\mathbf{R}$ .

## 2. Proof of Theorem 1.1

The proof of Theorem 1.1 is divided into the proof of the case of a trivial surface-knot  $F$  and the proof of the case of a general surface-link  $F$ . In the argument, the O2-handle pair  $(D \times I, D' \times I)$  is fixed in the 3-space  $\mathbf{R}[0]$  and consider normal forms of the core disks  $E, E'$  of the 2-handles  $E \times I, E' \times I$  in  $\mathbf{R}^4$ . To avoid the complexity of handling the intersection point  $q = E \cap E'$ , a sufficiently small boundary-collar  $n(\partial E')$  of  $E'$  is fixed in  $\mathbf{R}^3[0]$  and consider a normal form of the disk  $E'_n = \text{cl}(E' \setminus n(\partial E'))$  in  $\mathbf{R}^4$  together with a normal form of  $E$ .

**Proof of Theorem 1.1 in the case of a trivial surface-link  $F$ .** Assume that the trivial surface-knot  $F$  is embedded standardly in  $\mathbf{R}^3[0]$  with a standard O2-handle pair  $(D \times I, D' \times I)$  on  $F$ . By [4], the disk union  $G = E \cup E'_n$  is deformed into a disk union  $G_1$  in the following form by an isotopy of  $\mathbf{R}^4$  keeping the boundary

$\partial G = \partial E \cup \partial E'_n$  (which is a trivial link in  $\mathbf{R}^3[0]$ ),  $n(\partial E')$  and  $F$  fixed:

$$G_1 \cap \mathbf{R}^3[t] = \begin{cases} \emptyset, & \text{for } t > 2, \\ \mathbf{d}'[t], & \text{for } t = 2, \\ o'[t], & \text{for } 1 < t < 2, \\ (\partial G \cup \ell \cup \mathbf{b}')[t], & \text{for } t = 1, \\ (\partial G \cup \ell)[t], & \text{for } 0 \leq t < 1, \\ \ell[t], & \text{for } -1 < t < 0, \\ (o \cup \mathbf{b})[t], & \text{for } t = -1, \\ o[t], & \text{for } -2 < t < -1, \\ \mathbf{d}[t], & \text{for } t = -2, \\ \emptyset, & \text{for } t < -2, \end{cases}$$

where the notations  $o, o'$  denote trivial links in  $\mathbf{R}^3$ , the notations  $\mathbf{d}, \mathbf{d}'$  denote disjoint disk systems in  $\mathbf{R}^3$  bounded by  $o, o'$ , respectively, the notations  $\mathbf{b}, \mathbf{b}'$  denote disjoint band systems in  $\mathbf{R}^3$  spanning  $o, o'$ , respectively, and the notation  $\ell$  denotes a link in  $\mathbf{R}^3$ . To obtain this disk union  $G_1$ , start the argument of [4] with the assumption that the intersection  $G \cap \mathbf{R}^3[0]$  is a link  $\ell[0] \cup \partial G$  in  $\mathbf{R}^3[0]$  and a boundary-collar  $n(\partial G)$  of  $\partial G$  in  $G$  is in  $\mathbf{R}^3[0, c]$  so that

$$n(\partial G) \cap \mathbf{R}^3[t] = \partial G[t], \quad t \in [0, c]$$

for a small number  $c > 0$ , where  $\partial G$  is regarded to be in  $\mathbf{R}^3$  under the canonical identification  $\mathbf{R}^3[0] = \mathbf{R}^3$ . Then pull down a minimal point of  $G$  in  $\mathbf{R}^3(0, \infty)$  to  $\mathbf{R}^3(-\infty, 0)$  and pull up a maximal point of  $G$  in  $\mathbf{R}^3(-\infty, 0)$  to  $\mathbf{R}^3(0, \infty)$ . In these deformations, trivial components are increased in the intersection link  $G \cap \mathbf{R}^3[0]$ . After these preparations, do normalizations of  $G \cap \mathbf{R}^3[0, \infty)$  and  $G \cap \mathbf{R}^3(-\infty, 0]$  keeping  $G \cap \mathbf{R}^3[0]$  fixed. The band systems  $\mathbf{b}, \mathbf{b}'$  are made disjoint by band slide and band thinning and disjoint from  $\partial G$  by band deformation. Let  $G_1 = E \cup E'_n$ . The following notation is used.

**Notation.** The disk subsystems of the disk system  $\mathbf{d}$  belonging to  $E$  or  $E'_n$  are denoted by  $\mathbf{d}(E)$  or  $\mathbf{d}(E'_n)$ , respectively. The band subsystems of the band system  $\mathbf{b}$  belonging to  $E$  or  $E'_n$  are denoted by  $\mathbf{b}(E)$  or  $\mathbf{b}(E'_n)$ , respectively.

A next deformation of  $G_1$  is to change the level of the band system  $\mathbf{b}(E)[-1]$  into  $\mathbf{b}(E)[1]$  and the level of the disk system  $\mathbf{d}(E)[-2]$  into  $\mathbf{d}(E)[0.5]$ . To do so, it is observed that in  $\mathbf{R}^3$ , the boundary  $\partial G$  and the band system  $\mathbf{b}(E'_n)$  meet the disk system  $\mathbf{d}(E)$  in finite interior points and in finite interior simple arcs, respectively. For a point  $x \in \mathbf{d}(E) \cap \partial G$ , find a point  $y \in \partial \mathbf{d}(E) \setminus \partial E$  and a simple arc  $\alpha$  from  $x$  to  $y$  in  $\mathbf{d}(E)$  which does not meet the band systems  $\mathbf{b}, \mathbf{b}'$  by band slide and band

thinning. Let  $n(\alpha)$  be a disk neighborhood of  $\alpha$  in  $\mathbf{d}(E)$ . Deform the disk system  $\mathbf{d}'(E)$  so that  $n(\alpha) \subset \mathbf{d}'(E)$ . Then the intersection  $e(\alpha) = n(\alpha)[-2, 2] \cap G_1$  is a disk in the interior of  $G_1$ . Let  $\tilde{e}(\alpha) = \text{cl}((\partial(n(\alpha)[-2, 2])) \setminus e(\alpha))$  be the complementary disk of the disk  $e(\alpha)$  in the 2-sphere  $\partial(n(\alpha)[-2, 2])$ . The disk union

$$\tilde{G}_1 = \text{cl}(G_1 \setminus e(\alpha)) \cup \tilde{e}(\alpha)$$

induces a normal form of the union of a deformed disk  $\tilde{E}$  of  $E$  and the disk  $E'_n$  with  $\partial\tilde{G}_1 = \partial G_1$ . Note that the disk  $\tilde{E}$  may meet with the surface  $F$  and the topological position of  $\tilde{E}$  in  $\tilde{G}_1$  may be changed from  $G_1$ , although the disk  $E' = E'_n \cup n(\partial E')$  is unchanged and the configuration of  $\tilde{G}_1$  is the same as  $G_1$ . Do this deformation for all points of the finite set  $\mathbf{d}(E) \cap \partial G$ . Further, for an arc  $\beta$  in the finite arc set  $\mathbf{d}(E) \cap \mathbf{b}(E'_n)$ , find a simple arc  $\alpha$  in  $\mathbf{d}(E)$  extending this arc  $\beta$  to a point  $y \in \partial\mathbf{d}(E) \setminus \partial E$  which does not meet the band systems  $\mathbf{b}, \mathbf{b}'$  by band slide and band thinning. For a disk neighborhood  $n(\alpha)$  in  $\mathbf{d}(E)$ , do the same deformation as above. Do this deformation for all arcs  $\beta$  in the finite arc set  $\mathbf{d}(E) \cap \mathbf{b}(E'_n)$ . Let  $\tilde{G}_1 = \tilde{E} \cup E'_n$  be the disk union obtained from  $G_1 = E \cup E'_n$  by all these deformations, which is in a normal form with the same configuration as  $G_1$  and we have

$$\mathbf{d}(\tilde{E}) \cap (\partial E \cup n(\partial E')) = \mathbf{d}(\tilde{E}) \cap \mathbf{b}(E'_n) = \emptyset$$

although the disk  $\tilde{E}$  may meet  $F$ . Now change the level of  $\mathbf{b}(\tilde{E})[-1]$  into  $\mathbf{b}(\tilde{E})[1]$  and the level of  $\mathbf{d}(\tilde{E})[-2]$  into  $\mathbf{d}(\tilde{E})[0.5]$ . The resulting disk union  $G_2 = \tilde{E} \cup E'_n$  is in the following form:

$$G_2 \cap \mathbf{R}^3[t] = \left\{ \begin{array}{ll} \emptyset, & \text{for } t > 2, \\ \mathbf{d}'[t], & \text{for } t = 2, \\ o'[t], & \text{for } 1 < t < 2, \\ (\partial G \cup \cup o(\tilde{E}) \cup \mathbf{b}(\tilde{E}) \cup \ell(E'_n) \cup \mathbf{b}') [t], & \text{for } t = 1, \\ (\partial G \cup o(\tilde{E}) \cup \ell(E'_n)) [t], & \text{for } 0.5 < t < 1, \\ (\partial G \cup \mathbf{d}(\tilde{E}) \cup \ell(E'_n)) [t], & \text{for } t = 0.5, \\ (\partial G \cup \ell(E'_n)) [t], & \text{for } 0 \leq t < 0.5, \\ \ell(E'_n) [t], & \text{for } -1 < t < 0, \\ (o(E'_n) \cup \mathbf{b}(E'_n)) [t], & \text{for } t = -1, \\ o(E'_n) [t], & \text{for } -2 < t < -1, \\ \mathbf{d}(E'_n) [t], & \text{for } t = -2, \\ \emptyset, & \text{for } t < -2. \end{array} \right.$$

In the configuration of the disk union  $G_2$ , the pair  $(\tilde{E} \times I, E' \times I)$  is an O2-handle pair on  $F$  and hence is equivalent to the original O2-handle pair  $(E \times I, E' \times I)$  on  $F$  by Common 2-handle property. Let  $G_2 = E \cup E'_n$ . A next deformation of  $G_2$  is to

change the level of the band system  $\mathbf{b}(E'_n)[-1]$  into  $\mathbf{b}(E'_n)[1]$  and the level of the disk system  $\mathbf{d}(E'_n)[-2]$  into  $\mathbf{d}(E'_n)[0.5]$ . To do so, for a point  $x \in \mathbf{d}(E'_n) \cap \partial G$ , find a point  $y \in \partial \mathbf{d}(E'_n) \setminus \partial E'_n$  and a simple arc  $\alpha$  from  $x$  to  $y$  in  $\mathbf{d}(E'_n)$  which does not meet the band systems  $\mathbf{b}, \mathbf{b}'$  by band slide and band thinning. For a disk neighborhood  $n(\alpha)$  of  $\alpha$  in  $\mathbf{d}(E'_n)$ , do a similar deformation on the disk  $E'_n$  as above. Namely, deform the disk system  $\mathbf{d}'(E'_n)$  so that  $n(\alpha) \subset \mathbf{d}'(E'_n)$ . Since the intersection  $e(\alpha) = n(\alpha)[-2, 2] \cap G_2$  is a disk in the interior of  $G_2$ , let  $\tilde{e}(\alpha) = \text{cl}((\partial(n(\alpha)[-2, 2])) \setminus e(\alpha))$  be the complementary disk of the disk  $e(\alpha)$  in the 2-sphere  $\partial(n(\alpha)[-2, 2])$ . The disk union

$$\tilde{G}_2 = \text{cl}(G_2 \setminus e(\alpha)) \cup \tilde{e}(\alpha)$$

induces a normal form of the union of the disk  $E$  and a deformed disk  $\tilde{E}'_n$  of  $E'_n$  with  $\partial \tilde{G}_2 = \partial G_2$ . Note that the disk  $\tilde{E}'_n$  may meet  $F$  and the topological position of  $\tilde{E}'_n$  in  $\tilde{G}_1$  may be changed from  $G_1$ , although the disk  $E$  is unchanged and the configuration of  $\tilde{G}_2$  is the same as  $G_2$ . Do this operation for all points of the finite set  $\mathbf{d}(E'_n) \cap \partial G$ . Let  $\tilde{G}_2 = E \cup \tilde{E}'_n$  be the disk union obtained from  $G_2$  by all these deformations. The disk union  $\tilde{G}_2 = E \cup \tilde{E}'_n$  is in a normal form with the same configuration as  $G_2$  and has

$$\mathbf{d}(\tilde{E}'_n) \cap (\partial E \cup n(\partial E')) = \emptyset,$$

although  $\tilde{E}'_n$  may meet  $F$ . Now change the level of the band system  $\mathbf{b}(\tilde{E}'_n)[-1]$  into  $\mathbf{b}(\tilde{E}'_n)[1]$  and the level of the disk system  $\mathbf{d}(\tilde{E}'_n)[-2]$  into  $\mathbf{d}(\tilde{E}'_n)[0.5]$ . The resulting disk union  $G_3 = E \cup \tilde{E}'_n$  is as follows:

$$G_3 \cap \mathbf{R}^3[t] = \begin{cases} \emptyset, & \text{for } t > 2, \\ \mathbf{d}'[t], & \text{for } t = 2, \\ \sigma'[t], & \text{for } 1 < t < 2, \\ (\partial G \cup \sigma \cup \mathbf{b} \cup \mathbf{b}')[t], & \text{for } t = 1, \\ (\partial G \cup \sigma)[t], & \text{for } 0.5 < t < 1, \\ (\partial G \cup \mathbf{d})[t], & \text{for } t = 0.5, \\ (\partial G)[t], & \text{for } 0 \leq t < 0.5, \\ \emptyset, & \text{for } t < 0. \end{cases}$$

In the disk union  $G_3$ , the pair  $(E \times I, \tilde{E}' \times I)$  with  $\tilde{E}' = \tilde{E}'_n \cup n(\partial E')$  is an O2-handle pair on  $F$  and hence equivalent to the original O2-handle pair  $(E \times I, E' \times I)$  by Common 2-handle property. Let  $G_3 = E \cup \tilde{E}'_n$ . In the configuration of  $G_3$ , the pairs  $(D \times I, E' \times I)$  and  $(E \times I, D' \times I)$  are O2-handle pairs on  $F$ . Thus, by Common 2-handle property, the O2-handle pairs  $(D \times I, D' \times I)$  and  $(E \times I, E' \times I)$  on  $F$  are equivalent. This completes the proof of Theorem 1.1 in the case of a trivial surface-link  $F$ .

**Proof of Theorem 1.1 in the case of a general surface-link  $F$ .** For a general surface-link  $F$  in  $\mathbf{R}^4$  and O2-handle pairs  $(D \times I, D' \times I)$  and  $(E \times I, E' \times I)$ , let  $F(D \times I, D' \times I)$  be the surface-link obtained by surgery along  $(D \times I, D' \times I)$  (see [2]). Let  $F'$  be a trivial surface-knot in  $\mathbf{R}^4$  obtained from the surface-link  $F(D \times I, D' \times I)$  obtained by surgery along 1-handles  $h_j (j = 1, 2, \dots, s)$  embedded in a connected Seifert hypersurface  $W$  for  $F(D \times I, D' \times I)$  avoiding the intersection loops  $E \cap W, E' \cap W$  (cf. [1]). Then there is a trivial torus-knot  $T$  in  $\mathbf{R}^4$  such that the connected sum  $F' \# T$  is a trivial surface-knot in  $\mathbf{R}^4$  obtained from  $F$  by surgery along the 1-handles  $h_j (j = 1, 2, \dots, s)$  and  $(D \times I, D' \times I)$  is a standard O2-handle pair on  $F' \# T$  attached to the connected summand  $T$ . By construction, the pairs  $(D \times I, D' \times I)$  and  $(E \times I, E' \times I)$  are O2-handles on the connected sum  $F' \# T$  attached to the connected summand  $T$  whose defining 4-ball is disjoint from the “2-handles”  $h_j (j = 1, 2, \dots, s)$  on  $F' \# T$  attached to  $F'$ . Let  $\mathbf{h}$  be the core disk system  $D(h_j), (j = 1, 2, \dots, s)$  of the 2-handle system  $h_j (j = 1, 2, \dots, s)$  on  $F' \# T$  attached to  $F'$ . By the proof for the case of a trivial surface-link  $F$ , the O2-handle pair  $(E \times I, E' \times I)$  is equivalent to  $(D \times I, D' \times I)$  on  $F' \# T$ . To obtain such an equivalence without crossing the core disk system  $\mathbf{h}$ , the proof is revised as follows: A normal form of the disk union  $\bar{G} = G \cup \mathbf{h} = E \cup E'_n \cup \mathbf{h}$  can be thought of as the following disk union  $\bar{G}_1$ :

$$\bar{G}_1 \cap \mathbf{R}^3[t] = \left\{ \begin{array}{ll} \emptyset, & \text{for } t > 2, \\ (d'(\mathbf{h}) \cup \mathbf{d}')[t], & \text{for } t = 2, \\ (o'(\mathbf{h}) \cup o')[t], & \text{for } 1 < t < 2, \\ (\partial \bar{G} \cup \ell(\mathbf{h}) \cup b'(\mathbf{h}) \cup \ell \cup \mathbf{b}')[t] & \text{for } t = 1, \\ (\partial \bar{G} \cup \ell(\mathbf{h}) \cup \ell)[t], & \text{for } 0 \leq t < 1, \\ (\ell(\mathbf{h}) \cup \ell)[t], & \text{for } -1 < t < 0, \\ (o(\mathbf{h}) \cup b(\mathbf{h}) \cup o \cup \mathbf{b})[t], & \text{for } t = -1, \\ (o(\mathbf{h}) \cup o)[t], & \text{for } -2 < t < -1, \\ (d(\mathbf{h}) \cup \mathbf{d})[t], & \text{for } t = -2, \\ \emptyset, & \text{for } t < -2, \end{array} \right. ,$$

where in addition to the notations on  $G_1$ , the following notations are also added. Namely, the notations  $o(\mathbf{h}), o'(\mathbf{h})$  denote trivial links in  $\mathbf{R}^3$  coming from  $\mathbf{h}$ , the notations  $d(\mathbf{h}), d'(\mathbf{h})$  denote disjoint disk systems in  $\mathbf{R}^3$  bounded by  $o(\mathbf{h}), o'(\mathbf{h})$ , respectively, coming from  $\mathbf{h}$ , the notations  $b(\mathbf{h}), b'(\mathbf{h})$  denote disjoint band systems in  $\mathbf{R}^3$  spanning  $o(\mathbf{h}), o'(\mathbf{h})$ , respectively, and the notation  $\ell(\mathbf{h})$  denotes a link in  $\mathbf{R}^3$  coming from  $\mathbf{h}$ . The band systems  $\mathbf{b}, \mathbf{b}', b(\mathbf{h}), b'(\mathbf{h})$  are made disjoint by band slide and band thinning. In this normal form  $\bar{G}_1$ , the disk system  $\mathbf{h}$  can be taken as

$$\mathbf{h} \cap D \times I = \mathbf{h} \cap D' \times I = \emptyset,$$

because the defining 4-ball of the connected summand  $T$  in the connected sum  $F' \# T$

contains the union  $D \times I \cup D' \times I$  and is disjoint from the 2-handles  $h_j$  ( $j = 1, 2, \dots, s$ ). By a method similar to the process from  $G_1$  to  $G_2$ , we have a deformation  $\tilde{G}_1 = \tilde{E} \cup E'_n \cup \mathbf{h}$  of  $\bar{G}_1$  with the same configuration as  $\bar{G}_1$  such that

$$\mathbf{d}(\tilde{E}) \cap (\partial E \cup n(\partial E')) = \mathbf{d}(\tilde{E}) \cap \mathbf{b}(E'_n) = \mathbf{d}(\tilde{E}) \cap b(\mathbf{h}) = \emptyset,$$

although  $\tilde{E}$  may meet  $F' \# T$ . Now change the level of  $\mathbf{b}(\tilde{E})[-1]$  into  $\mathbf{b}(\tilde{E})[1]$  and the level of  $\mathbf{d}(\tilde{E})[-2]$  into  $\mathbf{d}(\tilde{E})[0.5]$ . Then the disk union  $\bar{G}_2 = \tilde{E} \cup E'_n \cup \mathbf{h}$  obtained from  $\tilde{G}_1$  is as follows:

$$\bar{G}_2 \cap \mathbf{R}^3[t] = \begin{cases} \emptyset, & \text{for } t > 2, \\ (d'(\mathbf{h}) \cup \mathbf{d}')[t], & \text{for } t = 2, \\ (o'(\mathbf{h}) \cup o')[t], & \text{for } 1 < t < 2, \\ (\partial \bar{G} \cup \ell(\mathbf{h}) \cup b'(\mathbf{h}) \cup o(\tilde{E}) \cup \mathbf{b}(\tilde{E}) \cup \ell(E') \cup \mathbf{b}')[t], & \text{for } t = 1, \\ (\partial \bar{G} \cup \ell(\mathbf{h}) \cup o(\tilde{E}) \cup \ell(E'_n))[t], & \text{for } 0.5 < t < 1, \\ (\partial \bar{G} \cup \ell(\mathbf{h}) \cup \mathbf{d}(\tilde{E}) \cup \ell(E'_n))[t], & \text{for } t = 0.5, \\ (\partial \bar{G} \cup \ell(\mathbf{h}) \cup \ell(E'_n))[t], & \text{for } 0 \leq t < 0.5, \\ (\ell(\mathbf{h}) \cup \ell(E'_n))[t], & \text{for } -1 < t < 0, \\ (o(\mathbf{h}) \cup b(\mathbf{h}) \cup o(E'_n) \cup \mathbf{b}(E'_n))[t], & \text{for } t = -1, \\ (o(\mathbf{h}) \cup o(E'_n))[t], & \text{for } -2 < t < -1, \\ (d(\mathbf{h}) \cup \mathbf{d}(E'_n))[t], & \text{for } t = -2, \\ \emptyset, & \text{for } t < -2. \end{cases}$$

In the configuration of  $\bar{G}_2$ , the pair  $(\tilde{E} \times I, E' \times I)$  is an O2-handle pair on  $F' \# T$  and hence equivalent to the O2-handle pair  $(E \times I, E' \times I)$  on  $F' \# T$  by Common 2-handle property. Let  $\bar{G}_2 = E \cup E'_n \cup \mathbf{h}$ . By a similar consideration from  $G_2$  to  $G_3$ , we have a deformation  $\tilde{G}_2 = E \cup \tilde{E}'_n \cup \mathbf{h}$  of  $\bar{G}_2$  with the same configuration as  $\bar{G}_2$  such that

$$\mathbf{d}(\tilde{E}'_n) \cap (\partial E \cup n(\partial E')) = \mathbf{d}(\tilde{E}'_n) \cap b(\mathbf{h}) = \emptyset,$$

although the disk  $\tilde{E}'_n$  may meet  $F' \# T$ . Now change the level of  $\mathbf{b}(\tilde{E}'_n)[-1]$  into  $\mathbf{b}(\tilde{E}'_n)[1]$  and the level of  $\mathbf{d}(\tilde{E}'_n)[-2]$  into  $\mathbf{d}(\tilde{E}'_n)[0.5]$ . Then the disk union  $\bar{G}_3 =$



$E \cup \tilde{E}'_n \cup \mathbf{h}$  obtained from  $\tilde{G}_2$  is as follows:

$$\tilde{G}_3 \cap \mathbf{R}^3[t] = \left\{ \begin{array}{ll} \emptyset, & \text{for } t > 2, \\ (d'(\mathbf{h}) \cup \mathbf{d}')[t], & \text{for } t = 2, \\ (o'(\mathbf{h}) \cup o')[t], & \text{for } 1 < t < 2, \\ (\partial\tilde{G} \cup \ell(\mathbf{h}) \cup b'(\mathbf{h}) \cup o \cup \mathbf{b}' \cup \mathbf{b})[t], & \text{for } t = 1, \\ (\partial\tilde{G} \cup \ell(\mathbf{h}) \cup o)[t], & \text{for } 0.5 < t < 1, \\ (\partial\tilde{G} \cup \ell(\mathbf{h}) \cup \mathbf{d})[t], & \text{for } t = 0.5, \\ (\partial\tilde{G} \cup \ell(\mathbf{h})) [t], & \text{for } 0 \leq t < 0.5, \\ \ell(\mathbf{h})[t], & \text{for } -1 < t < 0, \\ (o(\mathbf{h}) \cup b(\mathbf{h})) [t], & \text{for } t = -1, \\ o(\mathbf{h})[t], & \text{for } -2 < t < -1, \\ d(\mathbf{h})[t], & \text{for } t = -2, \\ \emptyset, & \text{for } t < 0. \end{array} \right.$$

In the configuration of  $G_3$ , the pair  $(E \times I, \tilde{E}' \times I)$  with  $\tilde{E}' = \tilde{E}'_n \cup n(\partial E')$  is an O2-handle pair on  $F' \# T$  and hence equivalent to the O2-handle pair  $(E \times I, E' \times I)$  on  $F' \# T$  by Common 2-handle property. Let  $G_3 = E \cup E'_n \cup \mathbf{h}$ . Since  $(D \times I, D' \times I)$  is in  $\mathbf{R}^3[0]$ , the disk system  $\mathbf{h}$  is disjoint from the O2-handle pair  $(D \times I, D' \times I)$ , although the disk system  $d'(\mathbf{h})[2]$  is isotopically deformed in  $\mathbf{R}^3[2]$  in  $\tilde{G}_3$ . Thus, in the configuration of  $G_3$ , the pairs  $(D \times I, E' \times I)$  and  $(E \times I, D' \times I)$  are O2-handle pairs on  $F' \# T$  and disjoint from  $\mathbf{h}$ . This means that the O2-handle pairs  $(D \times I, D' \times I)$  and  $(E \times I, E' \times I)$  on  $F' \# T$  are equivalent under 3-cell moves disjoint from the 2-handles  $h_j$  ( $j = 1, 2, \dots, s$ ) by Common 2-handle property. By the back surgery from  $F' \# T$  to  $F$  on the 2-handles  $h_j$  ( $j = 1, 2, \dots, s$ ) on  $F' \# T$ , this means that the O2-handle pairs  $(D \times I, D' \times I)$  and  $(E \times I, E' \times I)$  on  $F$  are equivalent under 3-cell moves disjoint from the 1-handles  $h_j$  ( $j = 1, 2, \dots, s$ ) on  $F$ . This completes the proof of Theorem 1.1 in the case of a general surface-link  $F$ .  $\square$

This completes the proof of Theorem 1.1.

### 3. Proof of Proposition 1.2

The *Finger move canceling* is the following operation to cancel a double point of an immersed disk  $D$  in  $\mathbf{R}^4$ .

**Finger Move Canceling.** Let  $D$  be an immersed disk in  $\mathbf{R}^4$  with  $\partial D$  embedded, and  $S$  a trivial  $S^2$ -knot in  $\mathbf{R}^4$  meeting the immersed disk  $D$  at just one point  $x$  different from the double points of  $D$ . Let  $y$  be a double point of  $D$ , and  $\alpha$  a simple arc in the disk  $D$  joining  $x$  and  $y$  not meeting the other double points of  $D$ . Let  $d_x$  be a disk neighborhood of  $x$  in  $D$ , and  $d_y$  a disk neighborhood  $d_y$  of  $y$  in the 2-sphere

$S$ , regarding the disks  $d_x$  and  $d_y$  as disk fibers of a normal disk bundle over  $D$  in  $\mathbf{R}^4$ . Let  $V_\alpha$  be a disk bundle over the arc  $\alpha$  in  $\mathbf{R}^4$  such that  $(D \cup S) \cap V_\alpha = d_x \cup \alpha \cup d_y$ . Then the immersed disk  $D_1$  with  $\partial D_1 = \partial D$  is constructed from the immersed disk  $D$  so that

$$D_1 = \text{cl}(D \setminus d_x) \cup \text{cl}(\partial V_\alpha \setminus (d_x \cup d_y)) \cup \text{cl}(S \setminus d_y).$$

The number of the double points of  $D_1$  is smaller than the number of the double points of  $D$  by 1.

The 2-sphere  $S$  in Finger Move Canceling is called a *canceling sphere*. If there is a canceling sphere  $S$ , then the immersed disk  $D$  is changed into an embedded disk  $D_*$  by Finger Move Canceling operations of parallel canceling spheres of  $S$ . By using Finger Move Canceling, the proof of Proposition 1.2 is done as follows:

**Proof of Proposition 1.2.** By assumption, the immersed O2-pair  $(D \times I, D' \times I)$  on a surface-link  $F$  in  $\mathbf{R}^4$  has  $D \times I$  as an immersed 2-handle on  $F$  and  $D' \times I$  as an embedded 2-handle on  $F$ . Let  $d'$  be a small disk neighborhood of a point  $p' \in D'$  in  $D'$ . By shrinking  $D' \times I$  as  $d' \times I$ , one finds a trivial  $S^2$ -knot  $S$  in  $\mathbf{R}^4$  such that  $S$  meets the immersed core disk  $D$  of  $D \times I$  at just one point  $x$  different from the double points of  $D$  and is disjoint from  $F$  and  $D' \times I$ . This 2-sphere  $S$  is used for a canceling sphere for the immersed disk  $D$ . By Finger Move Canceling, the immersed disk  $D$  is changed into an embedded disk  $D_*$ , meaning that the pair  $(D_* \times I, D' \times I)$  is an O2-handle pair on  $F$ . This completes the proof of Proposition 1.2.  $\square$

**Acknowledgements.** This work was partly supported by Osaka Central Advanced Mathematical Institute, Osaka Metropolitan University (MEXT Joint Usage/Research Center on Mathematics and Theoretical Physics JPMXP0619217849). The author thanks Junpei Yasuda (a graduate student at Osaka University under the adviser Seiichi Kamada) for asking a question on [2, Claim 3.2.1] that motivated to write this paper.

## references

- [1] F. Hosokawa and A. Kawauchi, Proposals for unknotted surfaces in four-space, Osaka J. Math. 16(1979), 233-248.
- [2] A. Kawauchi, Ribbonness of a stable-ribbon surface-link, I. A stably trivial surface-link, Topology and its Applications 301(2021), 107522 (16pages). arXiv:1804.02654

- [3] A. Kawauchi, Triviality of a surface-link with meridian-based free fundamental group. arXiv:1804.04269
- [4] A. Kawauchi, T. Shibuya and S. Suzuki, Descriptions on surfaces in four-space, I : Normal forms, Math. Sem. Notes, Kobe Univ. 10(1982), 75-125; II: Singularities and cross-sectional links, Math. Sem. Notes, Kobe Univ. 11(1983), 31-69.