

The Casson-Walker-Lescop invariant for branched cyclic covers of S^3 branched over a 2-bridge knot

Yasuyoshi Tsutsumi

Department of Mathematics, Osaka City University

Sugimoto 3-3-138, Sumiyoshi-ku, Osaka 558-8585, Japan

e-mail: ceu43500@nyc.odn.ne.jp

October 8th, 2004

Abstract

Let $D(b_1, \dots, b_{2m})$ be a 2-bridge knot, let $M_{b_1, \dots, b_{2m}}^r$ be the r -fold cyclic covering of S^3 branched over $D(b_1, \dots, b_{2m})$. We compute the Casson-Walker-Lescop invariant of $M_{b_1, \dots, b_{2m}}^r$ by using Lescop's formula and Chbili's method.

References

- [1] N. Chbili, *The Casson-Walker-Lescop invariant of periodic* , preprint.
- [2] C. Lescop, *A global surgery formula for the Casson-Walker invariant*, Annals of mathematics Studies **140**. Princeton University Press, Princeton, NJ (1996).
- [3] J. H. Przytycki and M. V. Sokolov, *Surgeries on periodic links and homology of periodic 3-manifold*, Math. Proc. Comb. Phil. Soc. **131**, 295–307 (2001).