

Research plan

Michio Yoshiwaki

1. As the final step of a review of basic facts on tilting theory in Grothendieck categories from the point of view of derived categories by using 1-tilting stalk complexes, I will study Grothendieck groups. Let \mathcal{A} be a locally noetherian Grothendieck category. A Grothendieck group $K_0(\mathcal{A})$ is defined as follows :

$$K_0(\mathcal{A}) := \bigoplus_{[X] \in \text{fg } \mathcal{A}/\cong} \mathbb{Z}[X]/I$$

where $\text{fg } \mathcal{A}$ is the category of finitely generated objects of \mathcal{A} and I is the subgroup generated by

$$\langle [X] - [Y] + [Z] \mid 0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0 \text{ exact} \rangle.$$

This definition is a natural generalization of Grothendieck group in the module categories. Let T be a 1-tilting stalk complex. Put $B = \text{End}_{\mathcal{A}}(T)$. I will first give an alternative proof of $K_0(\mathcal{A}) \cong K_0(B)$.

2. The second purpose is to generalize my results on tilting theory in Grothendieck categories. Here, we suppose that the derived category $\mathcal{D}(\mathcal{A})$ of a Grothendieck category \mathcal{A} is compactly generated. In this case, if $T \in \mathcal{A}$ is an n -tilting stalk complex, then it is an n -tilting object. But the converse is not true. A counter example due to D'Este is already known. I will observe under which conditions the converse is true. Based on this observation, I plan to prove basic facts on tilting theory in Grothendieck categories from the point of derived categories by using n -tilting stalk complexes.
3. The third purpose is to apply the results on tilting theory in Grothendieck categories to tilting sheaves. Namely, I will show that tilting sheaves are m -tilting stalk complexes for some m , and then plan to examine the differences between the definitions of tilting sheaves and of tilting stalk complexes.
4. I plan to study tilting modules. The study is related to the second purpose. Let A be a finite-dimensional algebra over an algebraically closed field and T a tilting module over A . Then the number of isomorphism classes of simple A -modules is equal to that of indecomposable direct summands of T . Now, there has been a problem whether this property may replace one of the conditions of a tilting module that there is an exact sequence: $0 \rightarrow A \rightarrow T_0 \rightarrow \cdots \rightarrow T_r \rightarrow 0$, where $T_0, \dots, T_r \in \text{add } T$. I will investigate under which conditions this is true. If T is a tilting module in the classical sense, this has been known to be true without conditions.