

これまでの研究成果のまとめ（英訳文）

田中利史

I am working on Geometric Topology. In particular, I am studying on low-dimensional topology using knot theory and the theory of 4-manifolds. Knot theory is the theory of the mathematical study of knots to investigate knotted simple closed curves in a 3-dimensional space. Though there are a lot of ways to study, I regard knot theory as a field of geometric topology. If we regard the theory of 4-manifolds as the theory for studying geometric properties of 4-manifolds, it is deeply concerned with knot theory. Now I explain my recent researches.

1. 1996-1999

I studied knot cobordism in terms of the properties of quasipositive knots. Particularly, I showed the existence of an infinite family of linearly independent knots with arbitrary gaps between 4-genera and topological 4-genera by showing important properties of quasipositive knots concerning knot cobordism. Maximal Thurston-Bennequin number is an invariant of knots and links derived from 3-dimensional contact geometry. I gave certain formula concerning Kauffman polynomials and Maximal Thurston-Bennequin numbers of positive knots and links. Moreover, I generalized the result to alternating knots and links.

2. 1999-2000

By studying the properties of the Jones polynomials of symmetric unions of knots, I gave an infinite family of counterexamples for a conjecture of T. Fiedler, given in 1999, about the cobordism invariance of the Jones polynomial of a knot. Furthermore, by developing the research, I gave a classification of 2-knots (embedded spheres in a 4-dimensional space) associated with symmetric unions.

3. 2002-2003

I researched on a linear representation of the fundamental group of the knot complement of a knot. The A -polynomial of a knot is an invariant derived from all $SL(2, C)$ -representations of the fundamental group of the knot complement. The problem that whether the knot which gives the trivial A -polynomial is the unknot was given in a paper of D. Cooper and D. Long. This problem had already been proved affirmatively for knots except satellite knots with trivial Alexander polynomials. I showed the existence of an infinite family of prime satellite knots with the trivial Alexander polynomials and the nontrivial A -polynomials.

4. 2003-2004

I studied on the colored Jones polynomials of links. Now it is considered to be important to find a formula for the colored Jones polynomial because it contributes to investigating the Kashaev-Murakami-Murakami volume conjecture (that states how to evaluate the Gromov invariant of a knot from its colored Jones polynomial). We obtained a formula for the N -colored Jones polynomial of a double of a knot K in terms of the colored Jones polynomial of K . It generalizes Masbaum's formula for K being the unknot. As a corollary, we showed that if the Kashaev-Murakami-Murakami volume conjecture for untwisted positive (or negative) doubles of knots is true, then the colored Jones polynomial detects the unknot.

5. 2004-2005

I investigated a relationship between knot cobordism and a clasper. A clasper is an important object in a study of finite type invariants. I showed that if a knot K is obtained from K' by using a graph clasper whose all components have the positive first Betti numbers, then there is a ribbon concordance from K' to K .

6. 2005-2006

I confirmed a question of E. Ferrand concerning a relation between HOMFLY polynomials and Kauffman polynomials. By collaborating with A. Stoimenow and D. Matei, we gave counterexamples for

a problem of Przytycki (Kirby's problem 1.91(2)) concerning the colored Jones polynomials and mutations of a knot. On the one hand, we have obtained a "gauge-theory free" proof for the existence of an exotic structure of a certain Casson handle using Rasmussen's invariant derived from Khovanov theory.

7. 2006-2007

We have studied on smooth structures on 4-manifolds. By using Rasmussen's invariant, we have shown the existence of an exotic smooth structure on any noncompact, connected smooth 4-submanifold of R^4 . Moreover we have obtained the same result for any noncompact, connected smooth 4-submanifold of $\sharp nCP^2$ for any positive integer n by using Donaldson theory.