

G -prime and G -primary G -ideals on G -schemes

Mitsuyasu Hashimoto

Joint with Mitsuhiro Miyazaki

Nagoya University

19 November, 2008

Notation

Notation 1

Throughout this talk,

- S : scheme
- G : an S -group scheme flat of finite type
- X : a G -scheme (i.e., an S -scheme with a left G -action)

We always assume that X is noetherian.

$\mu : G \times G \rightarrow G$ denotes the product, and $a : G \times X \rightarrow X$ denotes the action. Note that a is flat of finite type.

G -linearized \mathcal{O}_X -module

Definition 2 (Mumford)

A G -linearized \mathcal{O}_X -module (an equivariant (G, \mathcal{O}_X) -module) is a pair (\mathcal{M}, Φ) such that \mathcal{M} is an \mathcal{O}_X -module, and $\Phi : a^* \mathcal{M} \rightarrow p_2^* \mathcal{M}$ is an isomorphism of $\mathcal{O}_{G \times X}$ -modules such that

$$(\mu \times 1_X)^* \Phi : (\mu \times 1_X)^* a^* \mathcal{M} \rightarrow (\mu \times 1_X)^* p_2^* \mathcal{M}$$

agrees with

$$\begin{aligned} (\mu \times 1_X)^* a^* \mathcal{M} &\xrightarrow{\cong} (1_G \times a)^* a^* \mathcal{M} \xrightarrow{\Phi} (1_G \times a)^* p_2^* \mathcal{M} \\ &\xrightarrow{\cong} p_{23}^* a^* \mathcal{M} \xrightarrow{\Phi} p_{23}^* p_2^* \mathcal{M} \xrightarrow{\cong} (\mu \times 1_X)^* p_2^* \mathcal{M}, \end{aligned}$$

where $p_{23} : G \times G \times X \rightarrow G \times X$ is the projection.

Morphisms and submodules

Definition 3

A morphism $\varphi : (\mathcal{M}, \Phi) \rightarrow (\mathcal{N}, \Psi)$ of G -linearized \mathcal{O}_X -modules is a morphism $\varphi : \mathcal{M} \rightarrow \mathcal{N}$ such that $\Psi \circ (a^* \varphi) = (p_2^* \varphi) \circ \Phi$.

Definition 4

Let (\mathcal{M}, Φ) be a G -linearized \mathcal{O}_X -module. We say that \mathcal{N} is an equivariant (G, \mathcal{O}_X) -submodule of \mathcal{M} if \mathcal{N} is an \mathcal{O}_X -submodule of \mathcal{M} , and $\Phi(a^* \mathcal{N}) = p_2^* \mathcal{N}$ (note that a and p_2 are flat). If, moreover, $\mathcal{M} = \mathcal{O}_X$, then we say that \mathcal{N} is a G -ideal of \mathcal{O}_X .

The category $\text{Qch}(G, X)$

Theorem 5 (H—)

The category $\text{Qch}(G, X)$ of quasi-coherent G -linearized \mathcal{O}_X -modules is a locally noetherian abelian category, and (\mathcal{M}, Φ) is a noetherian object of $\text{Qch}(G, X)$ if and only if \mathcal{M} is coherent. The forgetful functor $F_X : \text{Qch}(G, X) \rightarrow \text{Qch}(X)$ given by $(\mathcal{M}, \Phi) \mapsto \mathcal{M}$ is faithful exact, and admits a right adjoint.

If it is convenient and there is no danger, we omit the Φ of (\mathcal{M}, Φ) , and we say that \mathcal{M} is in $\text{Qch}(G, X)$.

Operations on $\text{Qch}(G, X)$

Let $\mathcal{M}, \mathcal{N}, \mathcal{L}$ be in $\text{Qch}(G, X)$, \mathcal{I} be a G -ideal, and $\mathcal{M}_1, \mathcal{M}_2, \mathcal{M}_3$, and \mathcal{M}_λ be quasi-coherent equivariant (G, \mathcal{O}_X) -submodules of \mathcal{M} . Let \mathcal{L} and \mathcal{M}_3 be coherent. Then the following modules have structures of quasi-coherent G -linearized \mathcal{O}_X -modules.

- $\underline{\text{Tor}}_i^{\mathcal{O}_X}(\mathcal{M}, \mathcal{N}), \underline{\text{Ext}}_{\mathcal{O}_X}^i(\mathcal{L}, \mathcal{M}),$
- $\underline{H}_{\mathcal{I}}^i(\mathcal{M}) \cong \varinjlim \underline{\text{Ext}}_{\mathcal{O}_X}^i(\mathcal{O}_X/\mathcal{I}^n, \mathcal{M}),$
- The Fitting ideal $\underline{\text{Fitt}}_j(\mathcal{L}),$
- $\mathcal{M}_1 \cap \mathcal{M}_2, \sum_{\lambda} \mathcal{M}_{\lambda}, \mathcal{I}\mathcal{M}_1,$
- $\mathcal{M}_1 : \mathcal{M}_3, \mathcal{M}_1 : \mathcal{I}, \dots$

The star operation

Let \mathcal{M} be in $\text{Qch}(G, X)$, and \mathfrak{m} be an \mathcal{O}_X -submodule of \mathcal{M} . The sum of all quasi-coherent equivariant (G, \mathcal{O}_X) -submodules of \mathcal{M} contained in \mathfrak{m} is denoted by \mathfrak{m}^* . \mathfrak{m}^* is the largest quasi-coherent equivariant (G, \mathcal{O}_X) -submodule of \mathcal{M} contained in \mathfrak{m} .

Remark 6

This notation goes back at least to Matijevic-Roberts paper in 1974.

Let $Y = V(\mathfrak{a})$ be a closed subscheme of X . Then $Y^* := V(\mathfrak{a}^*)$ is the smallest G -stable closed subscheme of X containing Y .

Some formulas

From now on, all ideals and G -ideals are required to be coherent. All modules and G -linearized modules are required to be quasi-coherent.

Lemma 7

Let \mathcal{M} be in $\text{Qch}(G, X)$, \mathfrak{m} , \mathfrak{n} , and \mathfrak{m}_λ be \mathcal{O}_X -submodules of \mathcal{M} , and \mathcal{N} be a coherent equivariant (G, \mathcal{O}_X) -submodule of \mathcal{M} . Let \mathcal{I} be a G -ideal of \mathcal{O}_X . Then we have:

- $(\bigcap_\lambda \mathfrak{m}_\lambda^*)^* = (\bigcap_\lambda \mathfrak{m}_\lambda)^*$
- $\mathfrak{m}^* \cap \mathfrak{n}^* = (\mathfrak{m} \cap \mathfrak{n})^*$
- $(\mathfrak{m} : \mathcal{N})^* = \mathfrak{m}^* : \mathcal{N}$
- $(\mathfrak{m} : \mathcal{I})^* = \mathfrak{m}^* : \mathcal{I}$

G -prime G -ideal

Lemma 8

Let \mathcal{P} be a G -ideal of \mathcal{O}_X . Then the following are equivalent.

- There exists some ideal \mathfrak{p} of \mathcal{O}_X such that \mathfrak{p} is prime (i.e., $V(\mathfrak{p})$ is integral) and $\mathfrak{p}^* = \mathcal{P}$.
- $\mathcal{P} \neq \mathcal{O}_X$, and if \mathcal{I} and \mathcal{J} are G -ideals of \mathcal{O}_X and $\mathcal{I}\mathcal{J} \subset \mathcal{P}$, then $\mathcal{I} \subset \mathcal{P}$ or $\mathcal{J} \subset \mathcal{P}$.

Definition 9

If the equivalent conditions in the lemma are satisfied, we say that \mathcal{P} is a **G -prime** G -ideal.

The G -radical

Definition 10

Let \mathcal{I} be a G -ideal of \mathcal{O}_X . Then $V_G(\mathcal{I})$ denotes the set of G -prime ideals containing \mathcal{I} . We set $\sqrt[G]{\mathcal{I}} := (\bigcap_{\mathcal{P} \in V_G(\mathcal{I})} \mathcal{P})^*$, and call $\sqrt[G]{\mathcal{I}}$ the G -radical of \mathcal{I} .

Lemma 11

Let \mathcal{I} , \mathcal{J} , and \mathcal{P} be G -ideals of \mathcal{O}_X . Then we have:

- $\mathcal{I} \subset \sqrt[G]{\mathcal{I}} \subset \sqrt{\mathcal{I}}$, $\sqrt[G]{\mathcal{I}} = \sqrt{\mathcal{I}}^*$
- If $\mathcal{I} \supset \mathcal{J}$, then $\sqrt[G]{\mathcal{I}} \supset \sqrt[G]{\mathcal{J}}$.
- $\sqrt[G]{\mathcal{I}\mathcal{J}} = \sqrt[G]{\mathcal{I} \cap \mathcal{J}} = \sqrt[G]{\mathcal{I}} \cap \sqrt[G]{\mathcal{J}}$.
- $\sqrt{\sqrt[G]{\mathcal{I}}} = \sqrt[G]{\mathcal{I}}$.
- If \mathcal{P} is a G -prime, then $\sqrt[G]{\mathcal{P}} = \mathcal{P}$.

G -radical G -ideal

Lemma 12

Let \mathcal{I} be a G -ideal of \mathcal{O}_X . Then the following are equivalent.

- $\mathcal{I} = \sqrt[\mathcal{G}]{\mathcal{I}}$
- \mathcal{I} is the intersection of finitely many G -prime G -ideals.
- There exists some ideal \mathfrak{a} of \mathcal{O}_X such that \mathfrak{a} is radical (i.e., $V(\mathfrak{a})$ is reduced), and $\mathfrak{a}^* = \mathcal{I}$.

Definition 13

If the equivalent conditions in the lemma are satisfied, then we say that \mathcal{I} is **G -radical**.

A G -prime G -ideal is G -radical.

G -primary submodules

From now on, until the end of the talk, let \mathcal{M} be a coherent G -linearized \mathcal{O}_X -module, and \mathcal{N} its coherent equivariant (G, \mathcal{O}_X) -submodule.

Definition 14

We say that \mathcal{N} is **G -primary** if $\mathcal{N} \neq \mathcal{M}$, and for any coherent equivariant (G, \mathcal{O}_X) -submodule \mathcal{L} of \mathcal{M} , either $\mathcal{N} : \mathcal{L} = \mathcal{O}_X$ or $\mathcal{N} : \mathcal{L} \subset \sqrt[\mathcal{G}]{\mathcal{N} : \mathcal{M}}$ holds.

If \mathcal{N} is G -primary, then $\mathcal{P} = \sqrt[\mathcal{G}]{\mathcal{N} : \mathcal{M}}$ is G -prime. In this case, we say that \mathcal{N} is \mathcal{P} - G -primary.

A criterion

Lemma 15

- For a prime ideal \mathfrak{p} of \mathcal{O}_X , \mathfrak{p}^* is G -prime.
- For a radical ideal \mathfrak{a} of \mathcal{O}_X , \mathfrak{a}^* is G -radical.
- If \mathfrak{n} is a \mathfrak{p} -primary \mathcal{O}_X -submodule of \mathcal{M} , then \mathfrak{n}^* is a \mathfrak{p}^* - G -primary submodule of \mathcal{M} .
- For a G -primary submodule \mathcal{N} of \mathcal{M} , there exists some primary \mathcal{O}_X -submodule \mathfrak{n} of \mathcal{M} such that $\mathfrak{n}^* = \mathcal{N}$.

G -primary decomposition

Definition 16

An expression

$$\mathcal{N} = \mathcal{M}_1 \cap \cdots \cap \mathcal{M}_r$$

is called a G -primary decomposition if this equation holds, and each \mathcal{M}_i is a G -primary submodule of \mathcal{M} . We say that the decomposition is **minimal** if $\mathcal{N} \neq \bigcap_{j \neq i} \mathcal{M}_j$ for any i , and $\sqrt[\mathcal{G}]{\mathcal{M}_i : \mathcal{M}}$ is distinct.

The existence

Proposition 17

\mathcal{N} has a minimal G -primary decomposition.

Proof.

Let

$$\mathcal{N} = \mathfrak{m}_1 \cap \cdots \cap \mathfrak{m}_r$$

be a usual primary decomposition. Then

$$\mathcal{N} = \mathcal{N}^* = (\mathfrak{m}_1 \cap \cdots \cap \mathfrak{m}_r)^* = \mathfrak{m}_1^* \cap \cdots \cap \mathfrak{m}_r^*$$

is a G -primary decomposition. We can make it minimal, as usual. \square

G -associated G -prime

Theorem 18

The set

$$\text{Ass}_G(\mathcal{M}/\mathcal{N}) = \{ \sqrt[G]{\mathcal{M}_i} : \mathcal{M} \mid i = 1, \dots, r \}$$

is independent of the choice of minimal G -primary decomposition

$$\mathcal{N} = \mathcal{M}_1 \cap \dots \cap \mathcal{M}_r,$$

and depends only on \mathcal{M}/\mathcal{N} .

We call an element of $\text{Ass}_G(\mathcal{M}/\mathcal{N})$ a G -associated G -prime. The set of minimal elements of $\text{Ass}_G(\mathcal{M}/\mathcal{N})$ is denoted by $\text{Min}_G(\mathcal{M}/\mathcal{N})$, and its element is called a G -minimal G -prime. An element of $\text{Ass}_G(\mathcal{M}/\mathcal{N}) \setminus \text{Min}_G(\mathcal{M}/\mathcal{N})$ is called a G -embedded G -prime.

G -primary and primary decomposition

Theorem 19

Let

$$\mathcal{N} = \mathcal{M}_1 \cap \cdots \cap \mathcal{M}_r$$

be a minimal G -primary decomposition and

$$\mathcal{M}_j = \mathfrak{m}_{j,1} \cap \cdots \cap \mathfrak{m}_{j,s_j}$$

a minimal primary decomposition. Then

$$\mathcal{N} = \bigcap_{i=1}^r (\mathfrak{m}_{i,1} \cap \cdots \cap \mathfrak{m}_{i,s_i})$$

is a minimal primary decomposition.

No embedded prime of G -primary submodule

Proposition 20

A G -primary submodule \mathcal{N} of \mathcal{M} does not have an embedded prime. For each minimal prime \mathfrak{p} of \mathcal{M}/\mathcal{N} , we have $\mathfrak{p}^* = \sqrt[G]{\mathcal{N} : \mathcal{M}}$.

Corollary 21

We have

$$\text{Ass}(\mathcal{M}/\mathcal{N}) = \coprod_{i=1}^s \text{Ass}(\mathcal{M}/\mathcal{M}_i) = \coprod_{\mathcal{P} \in \text{Ass}_G(\mathcal{M}/\mathcal{N})} \text{Ass}(\mathcal{O}_X/\mathcal{P})$$

and

$$\text{Ass}_G(\mathcal{M}/\mathcal{N}) = \{\mathfrak{p}^* \mid \mathfrak{p} \in \text{Ass}(\mathcal{M}/\mathcal{N})\}$$

Another corollary

Corollary 22

We have $\text{Ass}(\mathcal{M}/\mathcal{N}) = \text{Min}(\mathcal{M}/\mathcal{N})$ if and only if $\text{Ass}_G(\mathcal{M}/\mathcal{N}) = \text{Min}_G(\mathcal{M}/\mathcal{N})$.

Smooth groups

Lemma 23

Assume that G is S -smooth. If \mathfrak{a} is a radical ideal of \mathcal{O}_X , then \mathfrak{a}^* is also radical. In particular, any G -radical G -ideal is radical.

Corollary 24

Assume that G is S -smooth. If \mathcal{I} is a G -ideal of \mathcal{O}_X , then $\sqrt{\mathcal{I}} = \sqrt[G]{\mathcal{I}}$. In particular, $\sqrt{\mathcal{I}}$ is a G -radical G -ideal.

Groups with connected fibers

Lemma 25

Assume that $G \rightarrow S$ has connected fibers. If \mathfrak{q} is a primary ideal of \mathcal{O}_X , then \mathfrak{q}^* is also primary. In particular, a G -primary G -ideal is primary.

Corollary 26

Assume that $G \rightarrow S$ has connected fibers. If \mathcal{I} is a G -ideal, then a minimal G -primary decomposition of \mathcal{I} is also a minimal primary decomposition.

Smooth groups with connected fibers

Corollary 27

Assume that $G \rightarrow S$ is smooth with connected fibers. If \mathfrak{p} is a prime, then \mathfrak{p}^* is also a prime. Any G -prime G -ideal is a prime. For a G -ideal \mathcal{I} of \mathcal{O}_X , any associated prime of \mathcal{I} is a G -prime G -ideal.

The dimension of the fiber

Theorem 28

Let $\mathfrak{0}$ be G -primary in \mathcal{O}_X . Then the dimension of the fiber of $p_2 : G \times X \rightarrow X$ is constant.

G -primary implies equi-dimensional

Theorem 29

Let $\mathfrak{0}$ be G -primary in \mathcal{O}_X . If X has an affine open covering $(\text{Spec } A_i)$ such that each A_i is Hilbert, universally catenary, and for any minimal prime P of A_i , the heights of maximal ideals of A_i/P are the same (for example, X is of finite type over a field or \mathbb{Z}). Then the dimensions of the irreducible components of X are the same.

Remark 30

There is an example of $G = X$ such that the dimensions of the irreducible components are different. The red assumptions are necessary.

G -primary ideal is unmixed

Theorem 31

Let \mathcal{Q} be a G -primary G -ideal of \mathcal{O}_X . Let x and y be the generic points of irreducible components of $V(\mathcal{Q})$. Then $\dim \mathcal{M}_x = \dim \mathcal{M}_y$.

Matijevic–Roberts type theorem

Theorem 32

Let $y \in X$ and $Y = \bar{y}$. Let η be the generic point of an irreducible component of Y^* . Then:

- If \mathcal{M}_η is maximal Cohen–Macaulay (resp. of finite injective dimension, projective dimension m , $\dim - \text{depth} = n$, torsionless, reflexive, G -dimension g), then so is \mathcal{M}_y .
- If $\mathcal{O}_{X,\eta}$ is a complete intersection, then so is $\mathcal{O}_{X,y}$.
- If G is smooth and $\mathcal{O}_{X,\eta}$ is regular, then $\mathcal{O}_{X,y}$ is regular.
- Assume that G is smooth and X is a locally excellent \mathbb{F}_p -scheme. If $\mathcal{O}_{X,\eta}$ is weakly F -regular (resp. F -regular, F -rational), then so is $\mathcal{O}_{X,y}$.

A Corollary on graded rings

Consider the case $S = \text{Spec } \mathbb{Z}$, $G = \mathbb{G}_m^n$, and $X = \text{Spec } A$ is affine. Then A is a \mathbb{Z}^n -graded ring.

Corollary 33

Let A be a locally excellent \mathbb{Z}^n -graded \mathbb{F}_p -algebra. Let P be a prime ideal of A , and let P^* be the prime ideal generated by homogeneous elements of P . If A_{P^*} is weakly F -regular (resp. F -regular, F -rational), then so is A_P .

A history of Matijevic–Roberts type theorem

Theorem 32 for graded rings (i.e., the case that $S = \text{Spec } \mathbb{Z}$, $G = \mathbb{G}_m^n$, and X affine) (excluding (weak) F -regularity and F -rationality):

- Conjectured by Nagata (for the case $n = 1$, for Cohen–Macaulay property).
- Proved by Hochster–Ratliff, Matijevic–Roberts, Aoyama–Goto, Matijevic, Goto–Watanabe, Cavaliere–Niesi, Avramov–Achilles.

General case (again excluding (weak) F -regularity and F -rationality):

- The case that S is noetherian affine, and G is affine, smooth with connected fibers (H—)
- G is smooth with connected fibers (Ohtani - H—, unpublished)
- General case: Theorem 32

G -artinian G -schemes

Definition 34

X is said to be G -artinian if every G -prime of \mathcal{O}_X is a G -minimal prime of 0 .

Corollary 35

A G -artinian G -scheme is Cohen–Macaulay.

Thank you. This slide is available at Hashimoto's home page.