

Good filtrations and strong F -regularity of the ring of U_P -invariants

MITSUYASU HASHIMOTO

Graduate School of Mathematics, Nagoya University
Chikusa-ku, Nagoya 464-8602 JAPAN
hasimoto@math.nagoya-u.ac.jp

Abstract

Let k be an algebraically closed field of positive characteristic, G a reductive group over k , and V a finite dimensional G -module. Let P be a parabolic subgroup of G , and U_P its unipotent radical. We prove that if $S = \text{Sym } V$ has a good filtration, then S^{U_P} is strongly F -regular.

1. Introduction

Throughout this paper, k denotes an algebraically closed field, and G a reductive group over k . We fix a maximal torus T and a Borel subgroup B which contains T . We fix a base Δ of the root system Σ of G so that B is negative. For any weight $\lambda \in X(T)$, we denote the induced module $\text{ind}_B^G(\lambda)$ by $\nabla_G(\lambda)$. We denote the set of dominant weights by X^+ . For $\lambda \in X^+$, we call $\nabla_G(\lambda)$ the *dual Weyl module* of highest weight λ . Note that for $\lambda \in X(T)$, $\text{ind}_B^G(\lambda) \neq 0$ if and only if $\lambda \in X^+$ [Jan, (II.2.6)], and if this is the case, $\nabla_G(\lambda) = \text{ind}_B^G(\lambda)$ is finite dimensional [Jan, (II.2.1)]. We denote $\nabla_G(-w_0\lambda)^*$ by $\Delta_G(\lambda)$, and call it the *Weyl module* of highest weight λ , where w_0 is the longest element of the Weyl group of G .

2010 *Mathematics Subject Classification*. Primary 13A50; Secondary 13A35. Key Words and Phrases. good filtration, F -regular, invariant subring.

We say that a G -module W is *good* if $\text{Ext}_G^1(\Delta_G(\lambda), W) = 0$ for any $\lambda \in X^+$. A filtration $0 = W_0 \subset W_1 \subset W_2 \subset \cdots \subset W_r$ or $0 = W_0 \subset W_1 \subset W_2 \subset \cdots$ of W is called a *good filtration* of W if $\bigcup_i W_i = W$, and for any $i \geq 1$, $W_i/W_{i-1} \cong \nabla_G(\lambda(i))$ for some $\lambda(i) \in X^+$. A G -module W has a good filtration if and only if W is good and of countable dimension [Don1]. See also [Fr] and [Has1, (III.1.3.2)].

Let V be a finite dimensional G -module. Let P be a parabolic subgroup of G containing B , and U_P its unipotent radical. The objective of this paper is to prove the following.

Corollary 5.5 *Let k be of positive characteristic. Let V be a finite dimensional G -module, and assume that $S = \text{Sym } V$ is good as a G -module. Then S^{U_P} is a finitely generated strongly F -regular Gorenstein UFD.*

An F -finite Noetherian ring R of characteristic p is said to be strongly F -regular if for any nonzerodivisor a of R , there exists some $r > 0$ such that the $R^{(r)}$ -linear map $aF^r : R^{(r)} \rightarrow R$ ($x^{(r)} \mapsto ax^{p^r}$) is $R^{(r)}$ -split [HH1]. See (2.1) for the notation. A strongly F -regular F -finite ring is F -regular in the sense of Hochster–Huneke [HH2], and hence it is Cohen–Macaulay normal ([HH3, (4.2)], [Kun], and [Vel, (0.10)]).

Under the same assumption as in Corollary 5.5, it has been known that S^G is strongly F -regular [Has2]. This old result is a corollary to our Corollary 5.5, since T is linearly reductive and $S^G = S^B = (S^U)^T$ is a direct summand subring of S^U . Under the same assumption as in Corollary 5.5, it has been proved that S^U is F -pure [Has6]. An F -finite Noetherian ring R of characteristic p is said to be F -pure if the Frobenius map $F : R^{(1)} \rightarrow R$ splits as an $R^{(1)}$ -linear map. Almost by definition, an F -finite strongly F -regular ring is F -pure, and hence Corollary 5.5 (or Corollary 4.14) yields this old result, too.

Popov [Pop3] proved that if the characteristic of k is zero, G is a reductive group over k , and A is a finitely generated G -algebra, then A has rational singularities if and only if A^U does so. Corollary 5.5 (or Corollary 4.14) can be seen as a weak characteristic p version of one direction of this result. For a characteristic p result related to the other direction, see Corollary 3.9.

Section 2 is preliminaries. We review the Frobenius twisting of rings, modules, and representations. We also review the basics of F -singularities such as F -rationality and F -regularity.

In Section 3, we study the ring theoretic properties of the invariant subring $k[G]^U$ of the coordinate ring $k[G]$. The main results of this section are Lemma 3.8 and Corollary 3.9.

In Section 4, we state and prove our main result for $P = B$. In order to do so, we introduce the notion of G -strong F -regularity and G - F -purity. These notions have already appeared in [Has2] essentially. Our main theorem in the most general form can be stated using these words (Theorem 4.12). As in [Has2], Steinberg modules play important roles.

In Section 5, we generalize the main results in Section 4 to the case of general P . Donkin's results on U_P -invariants of good G -modules play an important role here.

In Section 6, we give some examples. The first one is the action associated with a finite quiver. The second one is a special case of the first, and is a determinantal variety studied by De Concini and Procesi [DP]. The third one is also an example of the first. It gives some new understandings on the study of Goto–Hayasaka–Kurano–Nakamura [GHKN]. It also has some relationships with Miyazaki's study [Miy].

In Section 7, we prove the following.

Theorem 7.11 *Let S be a scheme, G a reductive S -group acting trivially on a Noetherian S -scheme X . Let M be a locally free coherent (G, \mathcal{O}_X) -module. Then*

$$\text{Good}(\text{Sym } M) = \{x \in X \mid \text{Sym}(\kappa(x) \otimes_{\mathcal{O}_{X,x}} M_x) \\ \text{is a good } (\text{Spec } \kappa(x) \times_S G)\text{-module}\},$$

and $\text{Good}(\text{Sym } M)$ is Zariski open in X .

For a reductive group G over a field which is not linearly reductive, there is a finite dimensional G -module V such that $(\text{Sym } V)^G$ is not Cohen–Macaulay [Kem]. On the other hand, in characteristic zero, a reductive group G is linearly reductive, and Hochster and Roberts [HR] proved that $(\text{Sym } V)^G$ is Cohen–Macaulay for any finite dimensional G -module V . Later, Boutot proved that $(\text{Sym } V)^G$ has rational singularities [Bt]. In view of Corollary 5.5 and Theorem 7.11, it seems that the condition $\text{Sym } V$ being good is an appropriate condition to ensure that the good results in characteristic zero still holds.

Acknowledgement: The author is grateful to Professor S. Donkin, Professor S. Goto, and Professor V. L. Popov for valuable advice. Special thanks are due to Professor K.-i. Watanabe for warm encouragement.

2. Preliminaries

(2.1) Throughout this paper, p denotes a prime number. Let K be a perfect field of characteristic p .

For a K -space V and $e \in \mathbb{Z}$, we denote the abelian group V with the new K -space structure $\alpha \cdot v = \alpha^{p^{-e}}v$ by $V^{(e)}$, where the product of $\alpha^{p^{-e}}$ and v in the right hand side is given by the original K -space structure of V . An element of V , viewed as an element of $V^{(e)}$ is sometimes denoted by $v^{(e)}$ to avoid confusion. Thus we have $v^{(e)} + w^{(e)} = (v + w)^{(e)}$ and $\alpha v^{(e)} = (\alpha^{p^{-e}}v)^{(e)}$. If $f : V \rightarrow W$ is a K -linear map, then $f^{(e)} : V^{(e)} \rightarrow W^{(e)}$ given by $f^{(e)}(v^{(e)}) = w^{(e)}$ is a K -linear map again. Note that $(?)^{(e)}$ is an autoequivalence of the category of K -vector spaces.

If A is a K -algebra, then $A^{(e)}$ with the multiplicative structure of A is a K -algebra. So $a^{(e)}b^{(e)} = (ab)^{(e)}$ for $a, b \in A$. If M is an A -module, then $M^{(e)}$ is an $A^{(e)}$ -module by $a^{(e)}m^{(e)} = (am)^{(e)}$. For a K -algebra A and $r \geq 0$, the r th Frobenius map $F^r = F_A^r : A \rightarrow A$ is defined by $F^r(a) = a^{p^r}$. Then $F^r : A^{(r+e)} \rightarrow A^{(e)}$ is a K -algebra map for $e \in \mathbb{Z}$. Note that $F^r(a^{(r+e)}) = (a^{p^r})^{(e)}$. $F^r : A^{(r+e)} \rightarrow A^{(e)}$ is also written as $(F^r)^{(e)}$.

In commutative ring theory, $A^{(e)}$ is sometimes denoted by ${}^{-e}A$ or A^{p^e} .

(2.2) For a K -scheme X , the scheme X with the new K -scheme structure $X \xrightarrow{f} \text{Spec } K \xrightarrow{a(F_K^{-e})} \text{Spec } K$ is denoted by $X^{(e)}$, where f is the original structure map of X as a K -scheme. So for a K -algebra A , $\text{Spec } A^{(e)}$ is identified with $(\text{Spec } A)^{(e)}$. The Frobenius map $F^r : X \rightarrow X^{(r)}$ is a K -morphism. Note that $(?)^{(e)}$ is an autoequivalence of the category of K -schemes with the quasi-inverse $(?)^{(-e)}$, and it preserves the product. So the canonical map $(X \times Y)^{(e)} \rightarrow X^{(e)} \times Y^{(e)}$ is an isomorphism. If G is a K -group scheme, then with the product $G^{(e)} \times G^{(e)} \cong (G \times G)^{(e)} \xrightarrow{\mu^{(e)}} G^{(e)}$, $G^{(e)}$ is a K -group scheme, and $F^r : G^{(e)} \rightarrow G^{(e+r)}$ is a homomorphism of K -group schemes. If V is a G -module, then $V^{(e)}$ is a $G^{(e)}$ -module in a natural way. Thus $V^{(r)}$ is a G -module again for $r \geq 0$ via $F^r : G \rightarrow G^{(r)}$. If V has a basis v_1, \dots, v_n , $g \in G(K)$, and $gv_j = \sum_i c_{ij}v_i$, then $gv_j^{(r)} = \sum_i c_{ij}^{p^r}v_i^{(r)}$. If A is a G -algebra, then $A^{(r)}$ is a G -algebra again. If M is a (G, A) -module, then

$M^{(r)}$ is a $(G, A^{(r)})$ -module. See [Has2].

(2.3) Let A be an \mathbb{F}_p -algebra. We say that A is *F-finite* if A is a finite $A^{(1)}$ -module. An *F-finite* Noetherian K -algebra is excellent [Kun].

Let A be Noetherian. We denote by A° the set $A \setminus \bigcup_{P \in \text{Min } A} P$, where $\text{Min } A$ denotes the set of minimal primes of A . Let M be an A -module and N its submodule. We define

$$\text{Cl}_A(N, M) = N_M^* := \{x \in M \mid \exists c \in A^\circ \exists e_0 \geq 1 \forall e \geq e_0 \\ x \otimes c^{(-e)} \in M/N \otimes_A A^{(-e)} \text{ is zero}\},$$

and call it the *tight closure* of N in M . Note that $\text{Cl}_A(N, M)$ is an R -submodule of M containing N [HH2, Section 8]. We say that N is *tightly closed* in M if $\text{Cl}_A(N, M) = N$. For an ideal I of A , $\text{Cl}_A(I, A)$ is simply denoted by I^* . If $I^* = I$, then we say that I is tightly closed.

We say that A is *very strongly F-regular* if for any $a \in A^\circ$, there exists some $r \geq 1$ such that the $A^{(r)}$ -linear map $aF_A^r : A^{(r)} \rightarrow A$ has a splitting. That is, there is an $A^{(r)}$ -linear map $\Phi : A \rightarrow A^{(r)}$ such that $\Phi aF^r = \text{id}_{A^{(r)}}$ [HH3], [Has5]. We say that A is *strongly F-regular* if $\text{Cl}_A(N, M) = N$ for any A -module M and its submodule N [Hoc, p. 166]. We say that A is *weakly F-regular* if $I = I^*$ for any ideal I of A [HH2]. We say that A is *F-regular* if for any prime ideal P of A , A_P is weakly *F-regular* [HH2]. We say that A is *F-rational* if $I = I^*$ for any ideal I generated by $\text{ht } I$ elements, where $\text{ht } I$ denotes the height of I .

Lemma 2.4. *Let A be a Noetherian \mathbb{F}_p -algebra.*

- (i) *If A is very strongly F-regular, then it is strongly F-regular. The converse is true, if A is either local, F-finite, or essentially of finite type over an excellent local ring.*
- (ii) *If A is strongly F-regular, then it is F-regular. An F-regular ring is weakly F-regular. A weakly F-regular ring is F-rational.*
- (iii) *A pure subring of a strongly F-regular ring is strongly F-regular.*
- (iv) *An F-rational ring is normal.*
- (v) *An F-rational ring which is a homomorphic image of a Cohen–Macaulay ring is Cohen–Macaulay.*

(vi) *A locally excellent F -rational ring is Cohen–Macaulay.*

(vii) *If $A = \bigoplus_{i \geq 0} A_i$ is graded and A_0 is a field, and if A is weakly F -regular, then A is very strongly F -regular.*

(viii) *A Gorenstein F -rational ring is strongly F -regular.*

Proof. (i) is [Has5, (3.6), (3.9), (3.35)]. (ii) is [Has5, (3.7)], [HH2, (4.15)], and [HH3, (4.2)]. (iii) is [Has5, (3.17)]. (iv) and (v) are [HH3, (4.2)]. (vi) is [Vel, (0.10)].

(vii) is [LS, (4.3)], if the field A_0 is F -finite. We prove the general case. By [HH2, (4.15)], $A_{\mathfrak{m}}$ is weakly F -regular, where $\mathfrak{m} = \bigoplus_{i > 0} A_i$ is the irrelevant ideal. Let K be the perfect closure (the largest purely inseparable extension) of A_0 , and set $B := K \otimes_{A_0} A$. Then B is purely inseparable over A . It is easy to see that $B_{\mathfrak{m}} := B \otimes_A A_{\mathfrak{m}}$ is a local ring whose maximal ideal is $\mathfrak{m}B_{\mathfrak{m}}$. By [HH3, (6.17)], $B_{\mathfrak{m}}$ is weakly F -regular. By the proof of [LS, (4.3)], $B_{\mathfrak{m}}$ and B are strongly F -regular. By [Has5, (3.17)], A is strongly F -regular. As A is finitely generated over the field A_0 , A is very strongly F -regular by (i).

(viii) Let A be a Gorenstein F -rational ring. By [HH3, (4.2)], $A_{\mathfrak{m}}$ is Gorenstein F -rational for any maximal ideal \mathfrak{m} of A . If $A_{\mathfrak{m}}$ is strongly F -regular for any maximal ideal \mathfrak{m} of A , then A is strongly F -regular by [Has5, (3.6)]. Thus we may assume that (A, \mathfrak{m}) is local. Let (x_1, \dots, x_d) be a system of parameters of A . Then an element of $H_{\mathfrak{m}}^d(A)$ as the d th cohomology group of the modified Čech complex [BH, (3.5)] is of the form $a/(x_1 \cdots x_d)^t$ for some $t \geq 0$ and $a \in A$. This element is zero if and only if $a \in (x_1^t, \dots, x_d^t)$, by [BH, (10.3.20)]. So this element is in the tight closure $(0)_{H_{\mathfrak{m}}^d(A)}^*$ of 0 if and only if $a \in (x_1^t, \dots, x_d^t)^* = (x_1^t, \dots, x_d^t)$, and hence $(0)_{H_{\mathfrak{m}}^d(A)}^* = 0$. As A is Gorenstein, $H_{\mathfrak{m}}^d(A)$ is isomorphic to the injective hull $E_A(A/\mathfrak{m})$ of the residue field A/\mathfrak{m} . By [Has5, (3.6)], A is strongly F -regular. \square

(2.5) Let K be a field of characteristic zero, and A a K -algebra of finite type. We say that A is of *strongly F -regular type* if there is a finitely generated \mathbb{Z} -subalgebra R of A and a finitely generated flat R -algebra A_R such that $A \cong K \otimes_R A_R$, and for any maximal ideal \mathfrak{m} of R , $R/\mathfrak{m} \otimes_R A_R$ is strongly F -regular. See [Har, (2.5.1)].

3. The invariant subring $k[G]^U$

(3.1) Let the notation be as in the introduction. Let Λ be an abelian group. We say that $A = \bigoplus_{\lambda \in \Lambda} A_\lambda$ is a Λ -graded G -algebra if A is both a G -algebra and a Λ -graded k -algebra, and each A_λ is a G -submodule of A for $\lambda \in \Lambda$. This is the same as to say that A is a $G \times \text{Spec } k\Lambda$ -algebra, where $k\Lambda$ is the group algebra of Λ over k . It is a commutative cocommutative Hopf algebra with each $\lambda \in \Lambda$ group-like.

We say that a \mathbb{Z} -graded k -algebra $A = \bigoplus_{i \in \mathbb{Z}} A_i$ is *positively graded* if $A_i = 0$ for $i < 0$ and $k \cong A_0$.

(3.2) Let the notation be as in the introduction.

We need to review Popov–Grosshans filtration [Pop2], [Grs2].

Let us fix (until the end of this section) a function $h : X(T) \rightarrow \mathbb{Z}$ such that (i) $h(X^+) \subset \mathbb{N} = \{0, 1, \dots\}$; (ii) $h(\lambda) > h(\mu)$ whenever $\lambda > \mu$; (iii) $h(\chi) = 0$ for $\chi \in X(G)$. Such a function h exists [Grs2, Lemma 6].

Let V be a G -module. For a poset ideal π of X^+ , we define $O_\pi(V)$ to be the sum of all the G -submodules W of V such that W belongs to π , that is, if $\lambda \in X^+$ and $W_\lambda \neq 0$, then $\lambda \in \pi$. $O_\pi(V)$ is the biggest G -submodule of V belonging to π . We set $\pi(n) := h^{-1}(\{0, 1, \dots, n\})$ for $n \geq 0$ and $\pi(n) = \emptyset$ for $n < 0$. We also define $V\langle n \rangle := O_{\pi(n)}(V)$.

For a G -algebra A , $(A\langle n \rangle)$ is a filtration of A . That is, $1 \in A\langle 0 \rangle \subset A\langle 1 \rangle \subset \dots$, $\bigcup_n A\langle n \rangle = A$, and $A\langle n \rangle \cdot A\langle m \rangle \subset A\langle n + m \rangle$. The Rees ring $\mathcal{R}(A)$ of A is the subring $\bigoplus_n A\langle n \rangle t^n$ of $A[t]$. Letting G act on t trivially, $A[t]$ is a G -algebra, and $\mathcal{R}(A)$ is a G -subalgebra of $A[t]$. So the associated graded ring $\mathcal{G}(A) := \mathcal{R}(A)/t\mathcal{R}(A)$ is also a G -algebra.

We denote the opposite of U by U^+ .

Theorem 3.3 (Grosshans [Grs1, Theorem 16]). *Let A be a G -algebra which is good as a G -module. There is a G -algebra isomorphism $\Phi : \mathcal{G}(A) \rightarrow (A^{U^+} \otimes k[G]^U)^T$, where U acts right regularly on $k[G]$, T acts right regularly on $k[G]^U$ (because T normalizes U), and G acts left regularly on $k[G]^U$ and trivially on A^{U^+} . \square*

The direct product $G \times G$ acts on the coordinate ring $k[G]$ by

$$((g_1, g_2)f)(g) = f(g_1^{-1}gg_2) \quad (f \in k[G], g, g_1, g_2 \in G(k)).$$

In particular, $k[G]$ is a $G \times B$ -algebra. Taking the invariant subring by the subgroup $U = \{e\} \times U \subset G \times B$, $k[G]^U$ is a $G \times T$ -algebra, since T

normalizes U . Thus $k[G]^U = \bigoplus_{\lambda \in X(T)} k[G]_{\lambda}^U$ is an $X(T)$ -graded G -algebra. As a G -module,

$$k[G]_{\lambda}^U = \{f \in k[G] \mid f(gb) = \lambda(b)f(g)\} \cong ((-\lambda) \otimes k[G])^B = \text{ind}_B^G(-\lambda)$$

for $\lambda \in X(B) = X(T)$ by the definition of induction, see [Jan, (I.3.3)]. Thus we have:

Lemma 3.4. $k[G]^U \cong \bigoplus_{\lambda \in X^+} \nabla_G(\lambda) \boxtimes (-\lambda)$ as a $G \times T$ -module. It is an integral domain. \square

The converse is also true.

Lemma 3.5. Let A be a $G \times T$ -algebra such that $A \cong \bigoplus_{\lambda \in X^+} \nabla_G(\lambda) \boxtimes (-\lambda)$ as a $G \times T$ -module and that A^{U^+} is a domain, where $U^+ = U^+ \times \{e\} \subset G \times T$. Then $A \cong k[G]^U$ as a $G \times T$ -algebra.

Proof. Let $\varphi : X^+ \rightarrow X(T) \times X(T)$ be the semigroup homomorphism given by $\varphi(\lambda) = (\lambda, -\lambda)$. A^{U^+} is a $\varphi(X^+)$ -graded domain, and each homogeneous component $A_{\varphi\lambda}^{U^+} = \nabla_G(\lambda)^{U^+} \boxtimes (-\lambda)$ is one-dimensional. So by [Has3, Lemma 5.5], $A^{U^+} \cong k\varphi(X^+)$ as an $X(T) \times X(T)$ -graded k -algebra.

Set $G' = G \times T$, and $T' = T \times T$. Define $h' : X(T') \cong X(T) \times X(T) \rightarrow \mathbb{Z}$ by $h'(\lambda, \mu) = h(\lambda)$. For a G' -algebra A' , we have a filtration of A' from h' as in (3.2). We denote the associated graded algebra by $\mathcal{G}'(A')$.

It is easy to see that $A \cong \mathcal{G}'(A)$, and this is isomorphic to $B := (k\varphi(X^+) \otimes (k[G]^U \boxtimes k[T]))^{T'}$ by Theorem 3.3 (applied to G'). We define $\psi : k[G]^U \rightarrow B$ by

$$\psi(a \otimes t_{-\lambda}) = (t_{\lambda} \otimes t_{-\lambda}) \otimes (a \otimes t_{-\lambda}) \otimes t_{\lambda},$$

where $a \in \nabla_G(\lambda)$ and for $\mu \in X(T)$, t_{μ} is the element μ considered as a basis element of $kX(T) = k[T]$. We consider that $\nabla_G(\lambda) \boxtimes (-\lambda) \subset k[G]^U$. With respect to the left regular action, t_{λ} is of weight $-\lambda$. So ψ is a $G \times T$ -algebra isomorphism. Thus $A \cong k[G]^U$ as a $G \times T$ -algebra, as desired. \square

Assume that G is semisimple simply connected. Then by [Pop2], $X(B) \rightarrow \text{Pic}(G/B)$ ($\lambda \mapsto \mathcal{L}(\lambda)$) is an isomorphism, where $\mathcal{L}(\lambda) = \mathcal{L}_{G/B}(\lambda)$ is the G -linearized invertible sheaf on the flag variety G/B , associated to the B -module λ , see [Jan, (I.5.8)]. Thus we have

Lemma 3.6. *If G is semisimple and simply connected, then the Cox ring (the total coordinate ring, see [Cox], [EKW]) $\text{Cox}(G/B)$ is isomorphic to $\bigoplus_{\lambda \in X^+} \nabla_G(\lambda)$, as an $X(T)$ -graded G -module (that is, $G \times T$ -module), where both $H^0(G/B, \mathcal{L}(\lambda)) \subset \text{Cox}(G/B)$ and $\nabla_G(\lambda)$ are assigned degree $-\lambda$. The Cox ring $\text{Cox}(G/B)$ is also an integral domain, and hence isomorphic to $k[G]^U$ as an $X(T)$ -graded G -algebra.*

Proof. The first assertion follows from the fact that $H^0(G/B, \mathcal{L}(\lambda)) = \nabla_G(\lambda)$ for $\lambda \in X^+$ and $H^0(G/B, \mathcal{L}(\lambda)) = 0$ for $\lambda \in X(B) \setminus X^+$ [Jan, (II.2.6)].

Consider the $\mathcal{O}_{G/B}$ -algebra $\mathcal{S} := \text{Sym}(\mathcal{L}(\lambda_1) \oplus \cdots \oplus \mathcal{L}(\lambda_l))$, where $\lambda_1, \dots, \lambda_l$ are the fundamental dominant weights. Being a vector bundle over G/B , $\text{Spec } \mathcal{S}$ is integral. Hence $\text{Cox}(G/B) \cong \Gamma(G/B, \mathcal{S})$ is a domain.

The last assertion follows from Lemma 3.5. □

Lemma 3.7. *If G is semisimple and simply connected, then $k[G]^U$ is a UFD.*

Proof. This is a consequence of Lemma 3.6 and [EKW, Corollary 1.2].

There is another proof. Popov [Pop2] proved that $k[G]$ is a UFD. Moreover, G does not have a nontrivial character, since $G = [G, G]$, see [Hum, (29.5)]. It follows easily that $k[G]^\times = k^\times$ by [Ros, Theorem 3]. As U is unipotent, U does not have a nontrivial character. The lemma follows from Remark 3 after Proposition 2 of [Pop1]. See also [Has7, (4.31)]. □

By [Grs1, (2.1)], $k[G]^U$ is finitely generated. See also [RR] and [Grs2, Theorem 9].

By [Has3, Lemma 5.6] and Lemma 3.4, $k[G]^U$ is strongly F -regular in positive characteristic, and strongly F -regular type in characteristic zero. In any characteristic, $k[G]^U$ is Cohen–Macaulay normal.

In any characteristic, if G is semisimple simply connected, being a finitely generated Cohen–Macaulay UFD, $k[G]^U$ is Gorenstein [Mur].

Combining the observations above, we have:

Lemma 3.8. *$k[G]^U$ is finitely generated. It is strongly F -regular in positive characteristic, and strongly F -regular type in characteristic zero. If G is semisimple and simply connected, then $k[G]^U$ is a Gorenstein UFD.*

Corollary 3.9. *Let k be of positive characteristic, and A be a G -algebra which is good as a G -module. If A^U is finitely generated and strongly F -regular, then A is finitely generated and F -rational.*

Proof. This is proved similarly to [Pop3, Proposition 10] and [Grs2, Theorem 17].

As U and U^+ are conjugate, $A^{U^+} \cong A^U$, and it is finitely generated and strongly F -regular by assumption. Note that A is finitely generated [Grs2, Theorem 9].

Note that $k[G]^U$ is finitely generated and strongly F -regular by Lemma 3.8. So the tensor product $A^{U^+} \otimes k[G]^U$ is finitely generated and strongly F -regular by [Has3, (5.2)]. Thus its direct summand subring $(A^{U^+} \otimes k[G]^U)^T$ is also finitely generated and strongly F -regular [HH1, (3.1)]. By Theorem 3.3, $\mathcal{G}(A)$ is finitely generated and strongly F -regular, hence is F -rational ([HH1, (3.1)] and [HH3, (4.2)]). By [HM, (7.14)], A is F -rational. \square

4. The main result

Let the notation be as in the introduction. In this section, the characteristic of k is $p > 0$.

(4.1) For a G -module W and $r \geq 0$, $W^{(r)}$ denotes the r th Frobenius twist of W , see Section 2 and [Jan, (I.9.10)].

Let ρ denote the half sum of positive roots. For $r \geq 0$, let St_r denote the r th Steinberg module $\nabla_G((p^r - 1)\rho)$, if $(p^r - 1)\rho$ is a weight of G . Note that $(p^r - 1)\rho$ is a weight of G if p is odd or $[G, G]$ is simply connected.

The following lemma, which is the dual assertion of [Has2, Theorem 3], follows immediately from [Jan, (10.6)].

Lemma 4.2. *Let $(p^r - 1)\rho$ be a weight of G for any $r \geq 0$. Let V be a finite dimensional G -module. Then there exists some $r_0 \geq 1$ such that for any $r \geq r_0$ and any subquotient W of V , any nonzero (or equivalently, surjective) G -linear map $\varphi : St_r \otimes W \rightarrow St_r$ admits a G -linear map $\psi : St_r \rightarrow St_r \otimes W$ such that $\varphi\psi = \text{id}_{St_r}$.*

We set $\tilde{G} = \text{rad } G \times \Gamma$, where $\text{rad } G$ is the radical of G , and $\Gamma \rightarrow [G, G]$ is the universal covering of the derived subgroup $[G, G]$ of G . Note that there is a canonical surjective map $\tilde{G} \rightarrow G$, and hence any G -module (resp. G -algebra) is a \tilde{G} -module (resp. \tilde{G} -algebra) in a natural way. The restriction functor $\text{res}_{\tilde{G}}^G$ is full and faithful.

Let $S = \bigoplus_{i \geq 0} S_i$ be a positively graded finitely generated G -algebra which is an integral domain.

Assume first that $(p^r - 1)\rho$ is a weight of G for $r \geq 0$. We say that S is G -strongly F -regular if for any nonzero homogeneous element a of S^G , there exists some $r \geq 1$ such that the $(G, S^{(r)})$ -linear map

$$\text{id} \otimes aF^r : St_r \otimes S^{(r)} \rightarrow St_r \otimes S \quad (x \otimes s^{(r)} \mapsto x \otimes as^{p^r})$$

is a split mono. In general, we say that S is G -strongly F -regular if it is so as a \tilde{G} -algebra.

The following is essentially proved in [Has2]. We give a proof for completeness.

Lemma 4.3. *If S is a G -strongly F -regular positively graded finitely generated G -algebra domain, then S^G is strongly F -regular.*

Proof. We may assume that $G = \tilde{G}$. Let $A := S^G$.

As we assume that S is a finitely generated positively graded domain, A is a finitely generated positively graded domain, see [MFK, Appendix to Chapter 1, **A**]. Let a be a nonzero homogeneous element of A such that $A[1/a]$ is regular. Take $r \geq 1$ so that $\text{id} \otimes aF^r : St_r \otimes S^{(r)} \rightarrow St_r \otimes S$ is a split mono. Let $\Phi : St_r \otimes S \rightarrow St_r \otimes S^{(r)}$ be a $(G, S^{(r)})$ -linear map such that $\Phi \circ (\text{id} \otimes aF^r) = \text{id}$. Then consider the commutative diagram of $(G, A^{(r)})$ -modules

$$\begin{array}{ccccc} St_r \otimes A^{(r)} & \hookrightarrow & St_r \otimes S^{(r)} & \xrightarrow{\text{id}} & St_r \otimes S^{(r)} \\ \downarrow aF^r & & \downarrow aF^r & \nearrow \Phi & \\ St_r \otimes A & \hookrightarrow & St_r \otimes S & & \end{array}$$

Then applying the functor $\text{Hom}_G(St_r, ?)$ to this diagram, we get the commutative diagram of $A^{(r)}$ -modules

$$\begin{array}{ccccc} A^{(r)} & \xrightarrow{\text{id}} & A^{(r)} & \xrightarrow{\text{id}} & A^{(r)} \\ \downarrow aF^r & & \downarrow & \nearrow & \\ A & \longrightarrow & \text{Hom}_G(St_r, St_r \otimes S) & & \end{array}$$

see [Has2, Proposition 1, **5**]. This shows that the $A^{(r)}$ -linear map $aF^r : A^{(r)} \rightarrow A$ splits. By [HH1, (3.3)], A is strongly F -regular. \square

The following is also proved in [Has2] (see the proof of [Has2, Theorem 6]).

Theorem 4.4. *Let V be a finite dimensional G -module. If $S = \text{Sym } V$ has a good filtration (see the introduction for definition), then S is G -strongly F -regular.*

Lemma 4.5. *Let S be a G -strongly F -regular positively graded finitely generated G -algebra domain, and assume that there exists some $a \in S^G \setminus \{0\}$ such that $S[1/a]$ is strongly F -regular. Then S is strongly F -regular.*

Proof. We may assume that $G = \tilde{G}$. Let I be the radical ideal of S which defines the non-strongly F -regular locus of S . Such an ideal exists, see [HH1, (3.3)]. Then I is $G \times \mathbb{G}_m$ -stable, and hence $I \cap S^G$ is \mathbb{G}_m -stable. In other words, $I \cap S^G$ is a homogeneous ideal of S^G . By assumption, $0 \neq a \in I \cap S^G$. So $I \cap S^G$ contains a nonzero homogeneous element b . Take $r \geq 1$ so that $1 \otimes bF^r : St_r \otimes S^{(r)} \rightarrow St_r \otimes S$ has a spitting. Let x be any nonzero element of St_r . Then $x \otimes \text{id} : S^{(r)} \cong k \otimes S^{(r)} \rightarrow St_r \otimes S^{(r)}$ given by $s^{(r)} \mapsto x \otimes s^{(r)}$ is a split mono as an $S^{(r)}$ -linear map. Thus $(x \otimes \text{id}_S)(bF^r) = (\text{id} \otimes bF^r)(x \otimes \text{id}_{S^{(r)}})$ is a split mono as an $S^{(r)}$ -linear map, and hence so is $bF^r : S^{(r)} \rightarrow S$. By [HH1, (3.3)], S is strongly F -regular. \square

Let S be a finitely generated G -algebra. We say that S is G - F -pure if there exists some $r \geq 1$ such that $\text{id} \otimes F^r : St_r \otimes S^{(r)} \rightarrow St_r \otimes S$ splits as a $(G, S^{(r)})$ -linear map. Obviously, a G -strongly F -regular finitely generated positively graded G -algebra domain is G - F -pure. The following is essentially proved in [Has6].

Lemma 4.6. *Let S be a G - F -pure finitely generated G -algebra. Then S^G is F -pure.*

Proof. This is proved similarly to Lemma 4.3. See also [Has6]. \square

Lemma 4.7. *Let S and S' be a G - F -pure finitely generated G -algebras. Then the tensor product $S \otimes S'$ is G - F -pure.*

Proof. This is easy, and we omit the proof. \square

Lemma 4.8. *Let S be a G - F -pure finitely generated G -algebra, and assume that the $(G, S^{(r)})$ -linear map*

$$\text{id} \otimes F^r : St_r \otimes S^{(r)} \rightarrow St_r \otimes S$$

splits. Then the $(G, S^{(nr)})$ -linear map

$$\text{id} \otimes F^{nr} : St_{nr} \otimes S^{(nr)} \rightarrow St_{nr} \otimes S$$

splits for any $n \geq 0$.

Proof. Induction on n . The case that $n = 0$ is trivial. Assume that $n > 0$. Note that $St_{nr} \cong St_r \otimes St_{(n-1)r}^{(r)}$. So

$$\text{id} \otimes (F^{(n-1)r})^{(r)} : St_{nr} \otimes S^{(nr)} \rightarrow St_{nr} \otimes S^{(r)}$$

is identified with the map

$$\text{id} \otimes (\text{id} \otimes F^{(n-1)r})^{(r)} : St_r \otimes (St_{(n-1)r} \otimes S^{((n-1)r)})^{(r)} \rightarrow St_r \otimes (St_{(n-1)r} \otimes S)^{(r)},$$

and it has an $(G, S^{(nr)})$ -linear splitting by the induction assumption. On the other hand, $\text{id} \otimes F^r : St_{nr} \otimes S^{(r)} \rightarrow St_{nr} \otimes S$ splits by assumption, as $St_{nr} \cong St_r \otimes St_{(n-1)r}^{(r)}$. Thus the composite

$$St_{nr} \otimes S^{(nr)} \xrightarrow{\text{id} \otimes (F^{(n-1)r})^{(r)}} St_{nr} \otimes S^{(r)} \xrightarrow{\text{id} \otimes F^r} St_{nr} \otimes S,$$

which agrees with $\text{id} \otimes F^{nr}$, has a splitting, as desired. \square

Lemma 4.9. *Let $S = \bigoplus_{n \geq 0} S_n$ be a finitely generated positively graded G -algebra which is an integral domain. Then the following are equivalent.*

- 1** S is G -strongly F -regular.
- 2** S is $[G, G]$ -strongly F -regular.
- 3** S is Γ -strongly F -regular, where $\Gamma \rightarrow [G, G]$ is the universal covering.

Proof. The implications **1** \Rightarrow **2** \Leftrightarrow **3** is trivial. We prove the direction **3** \Rightarrow **1**. Replacing G by \tilde{G} if necessary, we may assume that $G = R \times \Gamma$, where R is a torus, and Γ is a semisimple and simply connected algebraic group. Let $a \in S^G$ be any nonzero homogeneous element. Then by assumption, the $(R, (S^{(r)})^\Gamma)$ -linear map

$$(aF^r)^* : \text{Hom}_{\Gamma, S^{(r)}}(St_r \otimes S, St_r \otimes S^{(r)}) \rightarrow \text{Hom}_{\Gamma, S^{(r)}}(St_r \otimes S^{(r)}, St_r \otimes S^{(r)})$$

is surjective. Taking the R -invariant,

$$(aF^r)^* : \text{Hom}_{G, S^{(r)}}(St_r \otimes S, St_r \otimes S^{(r)}) \rightarrow \text{Hom}_{G, S^{(r)}}(St_r \otimes S^{(r)}, St_r \otimes S^{(r)})$$

is still surjective, since R is linearly reductive. This is what we wanted to prove. \square

The following is proved similarly.

Lemma 4.10. *Let $S = \bigoplus_{n \geq 0} S_n$ be a finitely generated G -algebra. Then the following are equivalent.*

- 1 S is G - F -pure.
- 2 S is $[G, G]$ - F -pure.
- 3 S is Γ - F -pure, where $\Gamma \rightarrow [G, G]$ is the universal covering. □

Lemma 4.11. *Let G be semisimple and simply connected. Then $k[G]^U$ is G - F -pure.*

Proof. This is [Has6, Lemma 3]. □

The following is the main theorem of this paper.

Theorem 4.12. *Let $S = \bigoplus_{n \geq 0} S_n$ be a finitely generated positively graded G -algebra. Assume that*

- 1 S is F -rational and Gorenstein.
- 2 S is G - F -pure.

Then S is a G -strongly F -regular integral domain.

Proof. Note that S is normal [HH3, (4.2)]. As S is positively graded, S is an integral domain.

Replacing G by Γ , where $\Gamma \rightarrow [G, G]$ is the universal covering, we may assume that G is semisimple and simply connected, by Lemma 4.9 and Lemma 4.10.

As S is G - F -pure, there exists some $l \geq 1$ such that $\text{id} \otimes F^l : St_l \otimes S^{(l)} \rightarrow St_l \otimes S$ has a $(G, S^{(l)})$ -linear splitting $\psi : St_l \otimes S \rightarrow St_l \otimes S^{(l)}$.

Note that any graded (G, S) -module which is rank one free as an S -module is of the form $S(n)$, where $S(n)$ is S as a (G, S) -module, but the grading is given by $S(n)_i = S_{n+i}$. In fact, let $-n$ be the generating degree of the rank one free graded (G, S) -module, say M , then $M_{-n} \otimes S \rightarrow M$ is a (G, S) -isomorphism. As M_{-n} is trivial as a G -module (since G is semisimple), $M_{-n} \cong k(n)$ as a graded G -module. Thus $M \cong k(n) \otimes S \cong S(n)$.

Let a be the a -invariant of the Gorenstein positively graded ring S . Namely, $\omega_S \cong S(a)$ (as a graded (G, S) -module, see the last paragraph). Then

$$\mathrm{Hom}_{S^{(r)}}(S, S^{(r)}) \cong \mathrm{Hom}_{S^{(r)}}(S, (\omega_S)^{(r)}(-p^r a)) \cong \omega_S(-p^r a) \cong S((1-p^r)a)$$

for $r \geq 0$.

Let σ be any nonzero element of $\mathrm{Hom}_{S^{(1)}}(S, S^{(1)})_{(p-1)a} \cong S_0$. As $S_0 = k$ is G -trivial, $\sigma : S \rightarrow S^{(1)}$ is $(G, S^{(1)})$ -linear of degree $(p-1)a$.

For $r \geq 0$, let σ_r be the composite

$$S \xrightarrow{\sigma} S^{(1)} \xrightarrow{\sigma^{(1)}} S^{(2)} \xrightarrow{\sigma^{(2)}} \dots \xrightarrow{\sigma^{(r-1)}} S^{(r)}.$$

It is $(G, S^{(r)})$ -linear of degree $(p^r - 1)a$. Note that $\sigma_u = \sigma_{u-r}^{(r)} \sigma_r$ for $u \geq r$.

Hence by the composite map

$$(1) \quad Q_{r,u} : \mathrm{Hom}_{S^{(r)}}(S, S^{(r)}) \xrightarrow{\sigma_{u-r}^{(r)}} \mathrm{Hom}_{S^{(r)}}(S, \mathrm{Hom}_{S^{(u)}}(S^{(r)}, S^{(u)})) \\ \cong \mathrm{Hom}_{S^{(u)}}(S^{(r)} \otimes_{S^{(r)}} S, S^{(u)}) \cong \mathrm{Hom}_{S^{(u)}}(S, S^{(u)}),$$

the element σ_r is mapped to σ_u , where the first map $\sigma_{u-r}^{(r)}$ maps $f \in \mathrm{Hom}_{S^{(r)}}(S, S^{(r)})$ to the map $x \mapsto f(x) \cdot \sigma_{u-r}^{(r)}$. More precisely, we have $Q_{r,u}(f) = \sigma_{u-r}^{(r)} \circ f$.

By the induction on u , it is easy to see that $Q_{1,u}$ is an isomorphism, and σ_u is a generator of the rank one S -free module $\mathrm{Hom}_{S^{(u)}}(S, S^{(u)})$. It follows that $Q_{r,u}$ is an isomorphism for any $u \geq r$.

We continue the proof of Theorem 4.12. Take a nonzero homogeneous element b of $A = S^G$. It suffices to show that there exists some $u \geq 1$ such that $\mathrm{id} \otimes bF^u : St_u \otimes S^{(u)} \rightarrow St_u \otimes S$ splits as a $(G, S^{(u)})$ -linear map.

As S is F -rational Gorenstein, it is strongly F -regular by Lemma 2.4, (viii). So there exists some $r \geq 1$ such that

$$(bF_S^r)^* : \mathrm{Hom}_{S^{(r)}}(S, S^{(r)}) \rightarrow \mathrm{Hom}_{S^{(r)}}(S^{(r)}, S^{(r)})$$

given by $(bF_S^r)^*(\varphi) = \varphi bF_S^r$ is surjective.

Let V be the degree $-(p^r - 1)a - d$ component of S , where d is the degree of b . Note that $V \cong \mathrm{Hom}_{S^{(r)}}(S, S^{(r)})_{-d}$ is mapped onto $k \cong \mathrm{Hom}_{S^{(r)}}(S^{(r)}, S^{(r)})_0$ by $(bF_S^r)^*$. In particular, $-(p^r - 1)a - d \geq 0$. So $a \leq 0$. If $S \neq k$, then it is easy to see that $a < 0$.

By Lemma 4.2, there exists some $u_0 \geq 1$ such that for any $u \geq u_0$, for any subquotient W of V , and any G -linear nonzero map $f : St_u \otimes W \rightarrow St_u$, there exists some G -linear map $g : St_u \rightarrow St_u \otimes W$ such that $fg = \text{id}$. Take $u \geq u_0$ such that $u - r$ is divisible by l .

Now the diagram

$$\begin{array}{ccc} \text{Hom}_{S^{(r)}}(S, S^{(r)}) & \xrightarrow{(bF^r)^*} & \text{Hom}_{S^{(r)}}(S^{(r)}, S^{(r)}) \\ \cong \downarrow Q_{r,u} & & \cong \downarrow Q_{0,u-r}^{(r)} \\ \text{Hom}_{S^{(u)}}(S, S^{(u)}) & \xrightarrow{(bF^r)^*} & \text{Hom}_{S^{(u)}}(S^{(r)}, S^{(u)}) \end{array}$$

is commutative. So the bottom $(bF^r)^*$ is surjective. Let us consider the surjection

$$(bF^r)^* : W = ((bF^r)^*)^{-1}(k \cdot \sigma_{u-r}^{(r)}) \cap \text{Hom}_{S^{(u)}}(S, S^{(u)})_{ap^r(p^{u-r}-1)-d} \rightarrow k \cdot \sigma_{u-r}^{(r)}.$$

By definition, W is contained in the degree $ap^r(p^{u-r} - 1) - d$ component of $\text{Hom}_{S^{(u)}}(S, S^{(u)}) = S(-a(p^u - 1))$, which is isomorphic to V as a G -module. So W is isomorphic to a G -submodule of V .

Let $E := \text{Hom}(St_u, St_u)$. Then by the choice of u_0 and u , there exists some G -linear map $g_1 : E \rightarrow E \otimes W$ such that the composite

$$E \xrightarrow{g_1} E \otimes W \xrightarrow{1 \otimes (bF^r)^*} E \otimes (k \cdot \sigma_{u-r}^{(r)})$$

maps φ to $\varphi \otimes \sigma_{u-r}^{(r)}$.

We identify $E \otimes \text{Hom}_{S^{(u)}}(S, S^{(u)})$ by $\text{Hom}_{S^{(u)}}(St_u \otimes S, St_u \otimes S^{(u)})$ in a natural way. Similarly, $E \otimes \text{Hom}_{S^{(r)}}(S^{(r)}, S)$ is identified with $\text{Hom}_{S^{(r)}}(St_u \otimes S^{(r)}, St_u \otimes S)$, and so on.

Then letting $\nu := g_1(\text{id}_{St_u})$, the composite

$$St_u \otimes S^{(r)} \xrightarrow{\text{id} \otimes bF^r} St_u \otimes S \xrightarrow{\nu} St_u \otimes S^{(u)}$$

agrees with $\text{id} \otimes \sigma_{u-r}^{(r)}$.

Since S is G - F -pure, $u - r$ is a multiple of l , and $St_u \cong St_r \otimes St_{u-r}^{(r)}$, there exists some $(G, S^{(u)})$ -linear map $\Phi : St_u \otimes S^{(r)} \rightarrow St_u \otimes S^{(u)}$ such that $\Phi \circ (\text{id}_{St_u} \otimes F_S^{u-r}) = \text{id}$ by Lemma 4.8.

Viewing Φ as an element of $E \otimes \text{Hom}_{S^{(u)}}(S^{(r)}, S^{(u)})$, let

$$\beta \in E \otimes \text{Hom}_{S^{(r)}}(S^{(r)}, S^{(r)})$$

be the element $(\text{id}_E \otimes (Q_{r,u}^{(r)})^{-1})(\Phi)$. In other words, $\beta : St_u \otimes S^{(r)} \rightarrow St_u \otimes S^{(r)}$ is the unique map such that the composite

$$St_u \otimes S^{(r)} \xrightarrow{\beta} St_u \otimes S^{(r)} \xrightarrow{\text{id} \otimes \sigma_{u-r}^{(r)}} St_u \otimes S^{(u)}$$

is Φ .

Write $\beta = \sum_i \varphi_i \otimes a_i^{(r)}$, where $\varphi_i \in E$ and $a_i \in S$. Define $\beta' \in E \otimes \text{Hom}_S(S, S)$ by $\beta' = \sum_i \varphi_i \otimes a_i^{(r)}$. Then it is easy to check that $(\text{id} \otimes bF^r) \circ \beta = \beta' \circ (\text{id} \otimes bF^r)$ as maps $St_u \otimes S^{(r)} \rightarrow St_u \otimes S$.

Combining the observations above, the whole diagram of $(G, S^{(u)})$ -modules

$$\begin{array}{ccccc}
 St_u \otimes S^{(u)} & \xrightarrow{\text{id} \otimes F^{u-r}} & St_u \otimes S^{(r)} & & \\
 \downarrow \text{id} & \nearrow \Phi & \downarrow \beta & \searrow \text{id} \otimes bF^r & \\
 St_u \otimes S^{(u)} & \xleftarrow{\text{id} \otimes \sigma_{u-r}^{(r)}} & St_u \otimes S^{(r)} & & St_u \otimes S \\
 & \nwarrow \nu & \downarrow \text{id} \otimes bF^r & \swarrow \beta' & \\
 & & St_u \otimes S & &
 \end{array}$$

is commutative. So $\text{id} \otimes bF^u : St_u \otimes S^{(u)} \rightarrow St_u \otimes S$ has a $(G, S^{(u)})$ -linear splitting $\nu\beta'$. \square

Corollary 4.13. *Let S be as in Theorem 4.12. Then S^U is finitely generated and strongly F -regular.*

Proof. Finite generation is by [Grs2, Theorem 9].

We prove the strong F -regularity. We may assume that G is semisimple and simply connected. Then $k[G]^U$ is a strongly F -regular Gorenstein domain by Lemma 3.8. Hence the tensor product $S \otimes k[G]^U$ is also a strongly F -regular Gorenstein domain, see [Has3, Theorem 5.2]. As S is assumed to be G - F -pure and $k[G]^U$ is G - F -pure by Lemma 4.11, the tensor product $S \otimes k[G]^U$ is also G - F -pure by Lemma 4.7. Hence by the theorem, $S \otimes k[G]^U$ is G -strongly F -regular. It follows that $(S \otimes k[G]^U)^G$ is strongly F -regular by Lemma 4.3. As $S^U \cong (S \otimes k[G]^U)^G$ (see the proof of [Grs1, (1.2)]). See also [Dol, Lemma 4.1]), we are done. \square

Corollary 4.14. *Let V be a finite dimensional G -module, and assume that $S = \text{Sym } V$ has a good filtration as a G -module. Then S^U is finitely generated and strongly F -regular.*

Proof. Follows immediately from Corollary 4.13 and Theorem 4.4. \square

5. The unipotent radicals of parabolic subgroups

Let the notation be as in the introduction. Let I be a subset of Δ . Let $L = L_I$ be the corresponding Levi subgroup $C_G(\bigcap_{\alpha \in I} (\text{Ker } \alpha)^\circ)$, where $(?)^\circ$ denotes the identity component, and C_G denotes the centralizer. Let $P = P_I$ be the parabolic subgroup generated by B and L . Let U_P be the unipotent radical of P . Let $B_L := B \cap L$, and U_L the unipotent radical of B_L .

Here are two theorems due to Donkin.

Theorem 5.1 (Donkin [Don3, (1.2)]). *Let w_0 and w_L denote the longest elements of the Weyl groups of G and L , respectively. For $\lambda \in X^+$, we have $\nabla_G(\lambda)^{U_P} \cong \nabla_L(w_L w_0 \lambda)$ as L -modules.*

Theorem 5.2 (Donkin [Don3, (1.4)], [Don4, (3.9)]). *Let*

$$0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$$

be a short exact sequence of G -modules. If M_1 is good, then

$$0 \rightarrow M_1^{U_P} \rightarrow M_2^{U_P} \rightarrow M_3^{U_P} \rightarrow 0$$

is exact. In other words, if M is a good G -module, then $R^i(H^0(U_P, ?) \circ \text{res}_{U_P}^G)(M) = 0$ for $i > 0$.

From these two theorems, it follows immediately:

Lemma 5.3. *Let M be a good G -module. Then M^{U_P} is a good L -module.*

So we have:

Proposition 5.4. *Let k be of positive characteristic. Let S be a finitely generated positively graded G -algebra. Assume that S is Gorenstein F -rational, and G - F -pure. Then S^{U_P} is finitely generated and F -rational.*

Proof. By Lemma 5.3, S^{U_P} is good as an L -module. By Corollary 4.13, $(S^{U_P})^{U_L} \cong S^U$ is finitely generated and strongly F -regular. By Corollary 3.9, applied to the action of L on S^{U_P} , we have that S^{U_P} is finitely generated and F -rational. \square

Corollary 5.5. *Let k be of positive characteristic. Let V be a finite dimensional G -module, and assume that $S = \text{Sym } V$ is good as a G -module. Then S^{U_P} is a finitely generated strongly F -regular Gorenstein UFD.*

Proof. As U_P is unipotent, S^{U_P} is a UFD by Remark 3 after Proposition 2 of [Pop1].

On the other hand, S satisfies the assumption of Proposition 5.4 by Theorem 4.4. So by Proposition 5.4, S^{U_P} is finitely generated and F -rational. Being a finitely generated Cohen–Macaulay UFD, it is Gorenstein [Mur], and hence is strongly F -regular by Lemma 2.4, (viii). \square

Remark 5.6. Let k be of characteristic zero. The characteristic-zero counterpart of Proposition 5.4 is stated as follows: If S is a finitely generated G -algebra with rational singularities, then S^{U_P} is finitely generated with rational singularities. This is proved in the same line as Proposition 5.4. Note that S^U is finitely generated with rational singularities by [Pop3, Corollary 4, Theorem 6]. Then applying [Pop3, Corollary 4, Theorem 6] again to the action of L on S^{U_P} , S^{U_P} is finitely generated and has rational singularities, since $(S^{U_P})^{U_L} \cong S^U$ is so.

The characteristic-zero counterpart of Corollary 5.5 is stated as follows: If S is a finitely generated G -algebra with rational singularities and is a UFD, then S^{U_P} is a Gorenstein finitely generated UFD which is of strongly F -regular type. As we have already seen, S^{U_P} is finitely generated with rational singularities. S^{U_P} is a UFD by Remark 3 after Proposition 2 of [Pop1]. As S^{U_P} is also Cohen–Macaulay [KKMS, p. 50, Proposition], S^{U_P} is Gorenstein [Mur]. A Gorenstein finitely generated algebra with rational singularities is of strongly F -regular type, see [Har, (1.1), (5.2)].

6. Applications

The following is pointed out in the proof of [SvdB, (5.2.3)].

Lemma 6.1. *Let K be a field of characteristic zero, H be an extension of a finite group scheme by a torus over K , and A a finitely generated H -algebra of strongly F -regular type. Then A^H is of strongly F -regular type.*

Proof. Set $B = A^H$. Let \bar{K} be the algebraic closure of K . As can be seen easily, if $\bar{K} \otimes_K B$ is of strongly F -regular type, then so is B . Since $\bar{K} \otimes_K B \cong (\bar{K} \otimes_K A)^{\bar{K} \otimes_K H}$, replacing K by \bar{K} , we may assume that K is algebraically closed. Then H is an extension of a finite group Γ by a split torus \mathbb{G}_m^r for some r . As $A^H \cong (A^{\mathbb{G}_m^r})^\Gamma$, we may assume that H is either a split torus \mathbb{G}_m^r or a finite group Γ .

Now we can take a finitely generated \mathbb{Z} -subalgebra R of K and a finitely generated flat R -algebra A_R such that $K \otimes_R A_R \cong A$, and for any closed point x of $\text{Spec } R$, $\kappa(x) \otimes_R A_R$ is strongly F -regular. Extending R if necessary, we have an action of H_R on A_R which is extended to the action of H on A , where $H_R = (\mathbb{G}_m)_R^r$ or $H_R = \Gamma$. Extending R if necessary, we may assume that $n \in R^\times$, where n is the order of Γ , if $H = \Gamma$.

Now set $B_R := A_R^{H_R}$.

If $H = (\mathbb{G}_m)_R^r$, then B_R is the degree zero component of the \mathbb{Z}^r -graded finitely generated R -algebra A_R , and it is finitely generated, and is a direct summand subring of A_R .

If $H = \Gamma$, then $B_R \rightarrow A_R$ is an integral extension and B_R is finitely generated by [AM, (7.8)]. As $\rho : A_R \rightarrow B_R$ given by $\rho(a) = (1/n) \sum_{\gamma \in \Gamma} \gamma a$ is a splitting, B_R is a direct summand subring of A_R .

In either case, B_R is finitely generated over R , so extending R if necessary, we may assume that B_R is R -flat. Note that $B \cong K \otimes_R B_R$, since K is R -flat, and the invariance is compatible with a flat base change. Note also that $\kappa(x) \otimes_R B_R$ is a direct summand subring of $\kappa(x) \otimes_R A_R$, and $\kappa(x) \otimes_R A_R$ is strongly F -regular. Hence $\kappa(x) \otimes_R B_R$ is strongly F -regular by Lemma 2.4, (iii). This shows that $A^H = B$ is of strongly F -regular type. \square

The following is a refinement of [SvdB, (5.2.3)].

Corollary 6.2. *Let K be a field of characteristic zero, H an affine algebraic group scheme over K such that H° is reductive. Let S be a finitely generated H -algebra which has rational singularities and is a UFD. Then S^H is of strongly F -regular type.*

Proof. Let $H' := [H^\circ, H^\circ]$. Then $\bar{K} \otimes_K H'$ is semisimple, and does not have a nontrivial character. Thus $S^{H'}$ has rational singularities by Boutot's theorem [Bt] and is a UFD by [Has7, (4.28)]. So it is of strongly F -regular type by Hara [Har, (1.1), (5.2)]. As $(H/H')^\circ$ is a torus, $S^H = (S^{H'})^{H/H'}$ is of strongly F -regular type by Lemma 6.1. \square

Theorem 6.3. *Let k be an algebraically closed field, and $Q = (Q_0, Q_1, s, t)$ a finite quiver, where Q_0 is the set of vertices, Q_1 is the set of arrows, and s and t are the source and the target maps $Q_1 \rightarrow Q_0$, respectively. Let $d : Q_0 \rightarrow \mathbb{N}$ be a map. For $i \in Q_0$, set $M_i := k^{d(i)}$, and let H_i be any closed subgroup scheme of $GL(M_i)$ of the following:*

- (1) $GL(M_i), SL(M_i)$;

- (2) $Sp_{d(i)}$ (in this case, $d(i)$ is required to be even);
- (3) $SO_{d(i)}$ (in this case, the characteristic of k must not be two);
- (4) Levi subgroup of any of (1)–(3);
- (5) Derived subgroup of any of (1)–(4);
- (6) Unipotent radical of a parabolic subgroup of any of (1)–(5);
- (7) Any subgroup H_i of $GL(M_i)$ with a closed normal subgroup N_i of H_i such that N_i is any of (1)–(6), and H_i/N_i is a linearly reductive group scheme. In characteristic zero, we require that $(H_i/N_i)^\circ$ is a torus.

Set $H := \prod_{i \in Q_0} H_i$ and $M := \prod_{\alpha \in Q_1} \text{Hom}(M_{s(\alpha)}, M_{t(\alpha)})$. Then $(\text{Sym } M^*)^H$ is finitely generated, and strongly F -regular if the characteristic of k is positive, and strongly F -regular type if the characteristic of k is zero.

Proof. If H_i satisfies (7) and the corresponding N_i satisfies (x), where $1 \leq \mathbf{x} \leq 6$, then we say that H_i is of type (7, x).

Note that $\text{Sym } M^* \cong \bigotimes_{\alpha \in Q_1} \text{Sym}(M_{s(\alpha)} \otimes M_{t(\alpha)}^*)$.

First we prove that $\text{Sym } M^*$ has a good filtration as an H -module if each H_i is as in (1)–(5). To verify this, we only have to show that $\text{Sym}(M_{s(\alpha)} \otimes M_{t(\alpha)}^*)$ is a good H -module for each α , by Mathieu’s tensor product theorem [Mat]. This module is trivial as an H_i -module if $i \neq s(\alpha), t(\alpha)$. Thus it suffices to show that this is good as an $H_{s(\alpha)} \times H_{t(\alpha)}$ -module if $s(\alpha) \neq t(\alpha)$, and as an $H_{s(\alpha)}$ -module if $s(\alpha) = t(\alpha)$, see [Has2, Lemma 4]. By [Has2, Lemma 3, 3, 5, 6] and [Don2, (3.2.7), (3.4.3)], the assertion is true for $H_{s(\alpha)}, H_{t(\alpha)}$ of (1)–(3). By Mathieu’s theorem [Mat, Theorem 1], the groups of type (4) is also allowed. By [Don2, (3.2.7)] again, the groups of type (5) is also allowed. By [Has2, Theorem 6], the conclusion of the theorem holds this case.

We consider the general case. If H_i is of the form (1)–(5), then considering $N_i = U_i \subset B_i \subset H_i$, where B_i is a Borel subgroup of H_i and U_i its unipotent radical, B_i is a group of the form (7), and as the H_i -invariant and the B_i -invariant are the same thing for an H_i -module, we may replace H_i by B_i without changing the invariant subring. Hence in this case, we may assume that H_i is of the form (7, 6). Clearly, a group of the form (6) is also of the form (7, 6), letting $N_i = H_i$. So we may assume that each H_i is of type (7). If $(\text{Sym } M^*)^N$ is strongly F -regular (type), where $N = \prod_{i \in Q_0} N_i$, then

$(\mathrm{Sym} M^*)^H \cong ((\mathrm{Sym} M^*)^N)^{H/N}$ is also strongly F -regular in positive characteristic, since $H/N \cong \prod_{i \in Q_0} H_i/N_i$ is linearly reductive and $(\mathrm{Sym} M^*)^H$ is a direct summand subring of $(\mathrm{Sym} M^*)^N$. In characteristic zero, $(H/N)^\circ$ is a torus, and we can invoke Lemma 6.1. Thus we may assume that each H_i is of the form **(1)–(6)**. Then again by the argument above, we may assume that each H_i is of the form **(7,6)**. Again by the argument above, we may assume that each H_i is of the form **(6)**. Now suppose that $H_i \subset G_i \subset GL(M_i)$, and each G_i is of the form **(1)–(5)**, and H_i is the unipotent radical of the parabolic subgroup P_i of G_i . Then letting $G := \prod G_i$ and $P := \prod P_i$, $H = \prod H_i$ is the unipotent radical of the parabolic subgroup P of G . As $\mathrm{Sym} M^*$ has a good filtration as a G -module by the first paragraph, $(\mathrm{Sym} M^*)^H$ is finitely generated and strongly F -regular (type) by Corollary 5.5 and Remark 5.6. \square

This covers Example 1 and Example 2 of [Has2], except that we do not consider the case $p = 2$ here, if O_n or SO_n is involved. For example,

Example 6.4. Let $Q = 1 \rightarrow 2 \rightarrow 3$, $(d(1), d(2), d(3)) = (m, t, n)$, $H_1 = H_3 = \{e\}$, and $H_2 = GL_t$. Then $M = \mathrm{Hom}(M_2, M_3) \times \mathrm{Hom}(M_1, M_2)$, and $M \rightarrow M//H$ is identified with

$$\pi : M \rightarrow Y_t = \{f \in \mathrm{Hom}(M_1, M_3) \mid \mathrm{rank} f \leq t\},$$

where $\pi(\varphi, \psi) = \varphi\psi$ (De Concini–Procesi [DP]). Thus (the coordinate ring of) Y_t is strongly F -regular (type), as was proved by Hochster–Huneke [HH4, (7.14)] (F -regularity and strong F -regularity are equivalent for positively graded rings, see Lemma 2.4).

Next we consider an example which really requires a group of type **(7)** in Theorem 6.3.

Let K be a field, and $M = K^m$, $N = K^n$. Let $1 \leq s \leq n$, and $\underline{a} = (0 = a_0 < a_1 < \cdots < a_s = n)$ be an increasing sequence of integers. Let \mathfrak{G} , \mathfrak{S} , and \mathfrak{T} be disjoint subsets of $\{1, \dots, s\}$ such that $\mathfrak{G} \amalg \mathfrak{S} \amalg \mathfrak{T} = \{1, \dots, s\}$. Let

$$H = H(\underline{a}; \mathfrak{G}, \mathfrak{S}, \mathfrak{T}) := \left(\begin{array}{cccc} \boxed{H_1} & & & \\ & \boxed{H_2} & & * \\ & & \ddots & \\ & & & \boxed{H_s} \\ O & & & \end{array} \right) \subset GL_m(K) \cong GL(M),$$

where H_l is $GL_{a_l - a_{l-1}}$ if $l \in \mathfrak{G}$, $SL_{a_l - a_{l-1}}$ if $l \in \mathfrak{S}$, and $\{E_{a_l - a_{l-1}}\}$ if $l \in \mathfrak{T}$.

Let us consider the symmetric algebra $S = \text{Sym}(M \otimes N)$. It is a graded polynomial algebra over K with each variable degree one. Let e_1, \dots, e_m and f_1, \dots, f_n be the standard bases of $M = K^m$ and $N = K^n$, respectively. For sequences $1 \leq c_1, \dots, c_u \leq m$ and $1 \leq d_1, \dots, d_u \leq n$, we define $[c_1, \dots, c_u \mid d_1, \dots, d_u]$ to be the determinant $\det(e_{c_i} \otimes f_{d_j})_{1 \leq i, j \leq u}$. It is a minor of the matrix $(e_i \otimes f_j)$ up to sign, or zero. Let Σ be the set of minors

$$\begin{aligned} \{[c_1, \dots, c_u \mid d_1, \dots, d_u] \mid 1 \leq u \leq \min(m, n), \\ 1 \leq c_1 < \dots < c_u \leq m, 1 \leq d_1 < \dots < d_u \leq n\}. \end{aligned}$$

We say that $[c_1, \dots, c_u \mid d_1, \dots, d_u] \leq [c'_1, \dots, c'_v \mid d'_1, \dots, d'_v]$ if $u \geq v$, and $c'_i \geq c_i$ and $d'_i \geq d_i$ for $1 \leq i \leq v$. It is easy to see that Σ is a distributive lattice.

Set $\epsilon := \min \mathfrak{G}$. For $1 \leq l < \epsilon$, set

$$\Gamma_l := \{[1, \dots, a_l \mid d_1, \dots, d_{a_l}] \mid 1 \leq d_1 < \dots < d_{a_l} \leq n\}$$

if $l \in \mathfrak{G}$, and

$$\begin{aligned} \Gamma_l := \{[c_1, \dots, c_u \mid d_1, \dots, d_u] \mid a_{l-1} < u \leq a_l, \\ 1 \leq c_1 < \dots < c_u \leq a_l, c_t = t \ (t \leq a_{l-1}), 1 \leq d_1 < \dots < d_u \leq n\} \end{aligned}$$

if $l \in \mathfrak{I}$. Set $\Gamma = \bigcup_{l < \epsilon} \Gamma_l$. Note that Γ is a sublattice of Σ .

It is well-known that S is an ASL on Σ over K [BH, (7.2.7)]. For the definition of ASL, see [BH, (7.1)].

Lemma 6.5. *Let B be a graded ASL on a poset Ω over a field K . Let Ξ be a subset of Ω such that for any two incomparable elements $\xi, \eta \in \Xi$,*

$$(2) \quad \xi\eta = \sum c_i m_i$$

in S with each m_i in the right hand side being a monomial of Ξ divisible by an element ξ_i in Ξ smaller than both ξ and η . Then the subalgebra $K[\Xi]$ of B is a graded ASL on Ξ .

Proof. We may assume that m_i in the right hand side of (2) has the same degree as that of $\xi\eta$ for each ξ, η , and m_i . For a monomial $m = \prod_{\omega \in \Omega} \omega^{c(\omega)}$, the weight $w(m)$ of m is defined to be $\sum_{\omega} c(\omega) 3^{\text{coht}(\omega)}$, where $\text{coht}(\omega)$ is the maximum of the lengths of chains $\omega = \omega_0 < \omega_1 < \dots$ in Ω . Then

$w(mm') = w(m) + w(m')$, and for each i , $w(m_i) > w(\xi\eta)$ in (2). So each time we use (2) to rewrite a monomial, the weight goes up. On the other hand, there are only finitely many monomials of a given degree, this rewriting procedure will stop eventually, and we get a linear combination of standard monomials in Ξ . Now (H_2) condition in [BH, (7.1)] is clear, while (H_0) and (H_1) are trivial. \square

We call $K[\Xi]$ a *subASL* of B generated by Ξ if the assumption of the lemma is satisfied.

Theorem 6.6. *Let the notation be as above. Let H act on S via $h(m \otimes n) = h(m) \otimes n$. Set $A := S^H$. Then*

- (1) $A = K[\Gamma]$.
- (2) $K[\Gamma]$ is a subASL of $S = K[\Sigma]$ generated by Γ .
- (3) A is a Gorenstein UFD. It is strongly F -regular if the characteristic of K is positive, and is of strongly F -regular type if the characteristic of K is zero.

Proof. First we prove that A is strongly F -regular (type). To do so, we may assume that $K = k$ is algebraically closed. Let B^+ be the subgroup of upper triangular matrices in GL_m , and set $B_H^+ := B^+ \cap H$. Then it is easy to see that $A = S^{B_H^+}$.

Now let Q be the quiver $1 \rightarrow 2$, $d = (d(1), d(2)) = (m, n)$, $G_1 = B_H^+ \subset GL_m$, and $G_2 = \{e\}$. Let U_H^+ be the unipotent radical of B_H^+ . Then U_H^+ is the unipotent radical of an appropriate parabolic subgroup of GL_m , U_H^+ is normal in B_H^+ , and B_H^+/U_H^+ is a torus. Thus the assumption (7) of Theorem 6.3 is satisfied, and thus $A = S^{B_H^+}$ is strongly F -regular (type).

The assertion (2) is a consequence of the straightening relation of the ASL S . See [ABW] for details.

Assume that (1) is proved. Then by the definition of Γ , letting M' be the subspace of M spanned by $e_1, \dots, e_{a_{\epsilon-1}}$, $A = \text{Sym}(M' \otimes N)^{H'}$, where $H' = H \cap GL(M')$, by (1) again ($GL(M')$ is viewed as a subgroup of $GL(M)$ via $g'(e_i) = e_i$ for $i > a_{\epsilon-1}$). As H' is connected and $\bar{K} \otimes_K H'$ does not have a nontrivial character, A is a UFD by [Has7, (4.28)], where \bar{K} is the algebraic closure of K . So assuming (1), the assertion (3) is proved.

It remains to prove (1). It is easy to see that $\Gamma \subset A$. So it suffices to prove that $\dim_K A_d = \dim_K K[\Gamma]_d$ for each degree $d \geq 0$. To do so, we may assume that K is algebraically closed.

Let P^+ be the parabolic subgroup $H(\underline{a}; \{1, \dots, s\}, \emptyset, \emptyset)$ of GL_m , and U_{P^+} the unipotent radical of P^+ . If

$$0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$$

is a short exact sequence of good $GL(M) \times GL(N)$ -modules, then

$$(3) \quad 0 \rightarrow M_1^{U_{P^+}} \rightarrow M_2^{U_{P^+}} \rightarrow M_3^{U_{P^+}} \rightarrow 0$$

is an exact sequence of good P^+/U_{P^+} -modules by Lemma 5.3 and Theorem 5.2. Note that P^+/U_{P^+} is identified with $\prod_{l=1}^s GL_{a_l - a_{l-1}}$, and H/U_{P^+} is identified with its subgroup $\prod_{l=1}^s H_l$. As each H_l is either $GL_{a_l - a_{l-1}}$, $SL_{a_l - a_{l-1}}$, or trivial, it follows that a good P^+/U_{P^+} -module is also good as an H/U_{P^+} -module. Applying the invariance functor $(?)^{H/U_{P^+}}$ to (3),

$$0 \rightarrow M_1^H \rightarrow M_2^H \rightarrow M_3^H \rightarrow 0$$

is exact.

Now we employ the standard convention for $GL(M)$. Let T be the set of diagonal matrices in $G := GL(M) = GL_m$, and we identify $X(T)$ with \mathbb{Z}^m by the isomorphism

$$\mathbb{Z}^m \ni (\lambda_1, \lambda_2, \dots, \lambda_m) \mapsto \left(\begin{pmatrix} t_1 & & & \\ & t_2 & & \\ & & \ddots & \\ & & & t_m \end{pmatrix} \mapsto t^\lambda = t_1^{\lambda_1} t_2^{\lambda_2} \cdots t_m^{\lambda_m} \right) \in X(T).$$

We fix the base of the root system of $GL(M)$ so that the set of lower triangular matrices in $GL(M)$ is negative. Then the set of dominant weights $X_{GL(M)}^+$ is the set

$$\{\lambda = (\lambda_1, \dots, \lambda_m) \in X(T) \mid \lambda_1 \geq \dots \geq \lambda_m\}.$$

We use a similar convention for $GL(N)$. See [Jan, (II.1.21)] for more information on this convention.

For $\lambda \in X_{GL(M)}^+$, $\nabla_{GL(M)}(\lambda)^{U_{P^+}}$ is a single dual Weyl module by Theorem 5.1. But obviously, the highest weight of $\nabla_{GL(M)}(\lambda)^{U_{P^+}}$ is λ . Thus $\nabla_{GL(M)}(\lambda)^{U_{P^+}} \cong \nabla_{P^+/U_{P^+}}(\lambda)$. Now the following is easy to verify:

Lemma 6.7. For $\lambda = (\lambda_1, \dots, \lambda_m) \in X_{GL(M)}^+$,

$$\nabla_{GL(M)}(\lambda)^H \cong \begin{cases} \nabla_{GL_{a_1}}(\lambda(1)) \otimes \cdots \otimes \nabla_{GL_{a_s - a_{s-1}}}(\lambda(s)) & (\lambda \in \Theta) \\ 0 & (\text{otherwise}) \end{cases}$$

as P^+/H -modules, where $\lambda(l) := (\lambda_{a_{l-1}+1}, \dots, \lambda_{a_l})$ for each l , and Θ is the subset of $X_{GL(M)}^+$ consisting of sequences $\lambda = (\lambda_1, \dots, \lambda_m)$ such that $\lambda(l) = (0, 0, \dots, 0)$ for each $l \in \mathfrak{G}$, and $\lambda(l) = (t, t, \dots, t)$ for some $t \in \mathbb{Z}$ for each $l \in \mathfrak{S}$.

Let $r := \min(m, n)$, and set

$$\mathcal{P}(d) = \{\lambda = (\lambda_1, \dots, \lambda_r) \in \mathbb{Z}^r \mid \lambda_1 \geq \cdots \geq \lambda_r \geq 0, |\lambda| = d\},$$

where $|\lambda| = \lambda_1 + \lambda_2 + \cdots + \lambda_r$. We consider that

$$(\lambda_1, \dots, \lambda_r) = (\lambda_1, \dots, \lambda_r, 0, \dots, 0),$$

and $\mathcal{P}(d) \subset X_{GL(M)}^+$. Similarly, we also consider that $\mathcal{P}(d) \subset X_{GL(N)}^+$. By the Cauchy formula [ABW, (III.1.4)], S_d has a good filtration as a $GL(M) \times GL(N)$ -module whose associated graded object is

$$\bigoplus_{\lambda \in \mathcal{P}(d)} \nabla_{GL(M)}(\lambda) \boxtimes \nabla_{GL(N)}(\lambda).$$

Note that $\nabla_{GL(M)}(\lambda)$ is isomorphic to the Schur module $L_{\tilde{\lambda}}M$ in [ABW], where $\tilde{\lambda}$ is the transpose of λ . That is, $\tilde{\lambda} = (\tilde{\lambda}_1, \tilde{\lambda}_2, \dots)$ is given by $\tilde{\lambda}_i = \#\{j \geq 1 \mid \lambda_j \geq i\}$.

By Lemma 6.7, S_d^H has a filtration whose associated graded object is

$$\bigoplus_{\lambda \in \mathcal{P}(d) \cap \Theta} \nabla_{GL_{a_1}}(\lambda(1)) \otimes \cdots \otimes \nabla_{GL_{a_s - a_{s-1}}}(\lambda(s)) \boxtimes \nabla_{GL(N)}(\lambda).$$

In particular,

$$(4) \quad \dim S_d^H = \sum_{\lambda \in \mathcal{P}(d) \cap \Theta} \dim \nabla_{GL(N)}(\lambda) \prod_l \dim \nabla_{GL_{a_l - a_{l-1}}}(\lambda(l)).$$

Next we count the dimension of $K[\Gamma]_d$. This is the number of standard monomials of degree d in $K[\Gamma]$. For a standard monomial

$$v = \prod_{b=1}^{\alpha} [c_{b,1}, \dots, c_{b,\mu_b} \mid \tilde{d}_{b,1}, \dots, \tilde{d}_{b,\mu_b}]$$

(where $[c_{b,1}, \dots, c_{b,\mu_b} \mid d_{b,1}, \dots, d_{b,\mu_b}]$ increases when b increases) in Σ , we define $\mu(v) = (\mu_1, \dots, \mu_\alpha)$, and $\lambda(v)$ its transpose. Such a standard monomial v of Γ of degree d exists if and only if $\lambda(v) \in \Theta \cap \mathcal{P}(d)$.

For a standard monomial v of Σ such that $\lambda(v) = \lambda \in \mathcal{P}(d) \cap \Theta$, v is a monomial of Γ if and only if the following condition holds. For each $1 \leq b \leq \lambda_1$, $1 \leq l \leq s$, and each $a_{l-1} < i \leq a_l$, it holds $a_{l-1} < c_{s,i} \leq a_l$. The number of such monomials agrees with $\dim \nabla_{GL(N)}(\lambda) \prod_l \dim \nabla_{GL_{a_l - a_{l-1}}}(\lambda(l))$, as can be seen easily from the standard basis theorem [ABW, (II.2.16)]. So $\dim_K K[\Gamma]$ agrees with the right hand side of (4), and we have $\dim_K A_d = \dim_K S_d^H = \dim_K K[\Gamma]_d$, as desired. \square

Remark 6.8. The case that $s = 2$, $a_1 = l$, $\mathfrak{G} = \emptyset$, $\mathfrak{S} = \{2\}$, and $\mathfrak{T} = \{1\}$ is studied by Goto–Hayasaka–Kurano–Nakamura [GHKN]. Gorenstein property and factoriality are proved there for this case. The case that $s = m$, $a_l = l$ ($l = 1, \dots, m$), $\mathfrak{G} = \mathfrak{S} = \emptyset$, and $\mathfrak{T} = \{1, \dots, m\}$ is a very special case of the study of Miyazaki [Miy].

7. Openness of good locus

(7.1) Let R be a Noetherian commutative ring, and G a split reductive group over R . We fix a split maximal torus T of G whose embedding into G is defined over \mathbb{Z} . We fix a base Δ of the root system, and let B be the negative Borel subgroup. For a dominant weight λ , the dual Weyl module $\nabla_G(\lambda)$ is defined to be $\text{ind}_B^G(\lambda)$, and the Weyl module $\Delta_G(\lambda)$ is defined to be $\nabla_G(-w_0\lambda)^*$.

A G -module M is said to be *good* if $\text{Ext}_G^1(\Delta_G(\lambda), M) = 0$ for any $\lambda \in X^+$, where X^+ is the set of dominant weights, see [Has1, (III.2.3.8)].

Lemma 7.2. *The notion of goodness of a G -module M is independent of the choice of T or Δ , and depends only on M .*

Proof. Let T' and Δ' be another choice of a split maximal torus defined over \mathbb{Z} and a base of the root system (with respect to T'). Let B' be the corresponding negative Borel subgroup.

Assume that R is an algebraically closed field. Then there exists some $g \in G(R)$ such that $gBg^{-1} = B'$. So $\text{ind}_B^G \lambda \cong \text{ind}_{B'}^G(\lambda')$ for any $\lambda \in X(B)$, where λ' is the composite

$$B' \xrightarrow{b' \mapsto g^{-1}b'g} B \xrightarrow{\lambda} \mathbb{G}_m.$$

So this case is clear.

When R is a field, then a G -module M is good if and only if $\bar{R} \otimes_R M$ is so as an $\bar{R} \otimes_R G$ -module, and this notion is independent of the choice of B , where \bar{R} is the algebraic closure of R .

Now consider the general case. If M is R -finite R -projective, then the assertion follows from [Has1, (III.4.1.8)] and the discussion above. If M is general, then M is good if and only if there exists some filtration

$$0 = M_0 \subset M_1 \subset M_2 \subset \dots$$

of M such that $\bigcup_i M_i = M$, and for each $i \geq 1$, $M_i/M_{i-1} \cong N_i \otimes V_i$ for some R -finite R -projective good G -module N_i and an R -module V_i . Indeed, the only if part is [Has1, (III.2.3.8)], while the if part is a consequence of the goodness of $N_i \otimes V_i$, see [Has1, (III.4.1.8)]. This notion is independent of the choice of T or Δ , and we are done. \square

Note that if $R \rightarrow R'$ is a Noetherian R -algebra, then an $R' \otimes_R G$ -module M' is good if and only if it is so as a G -module. This comes from the isomorphism

$$\mathrm{Ext}_G^i(\Delta_G(\lambda), M') \cong \mathrm{Ext}_{R' \otimes_R G}^i(\Delta_{R' \otimes_R G}(\lambda), M').$$

If M is a good G -module, and R' is R -flat or M is R -finite R -projective, then $R' \otimes_R M$ is a good $R' \otimes_R G$ -module by [Has1, (I.3.6.20)] and [Has1, (III.1.4.8)], see [Has1, (III.2.3.15)]. If M is good and V is a flat R -module, then $M \otimes V$ is good. This follows from the canonical isomorphism

$$\mathrm{Ext}_G^i(\Delta_G(\lambda), M \otimes V) \cong \mathrm{Ext}_G^i(\Delta_G(\lambda), M) \otimes V,$$

see [Has1, (I.3.6.16)].

If R' is faithfully flat over R and $R' \otimes_R M$ is good, then M is good by [Has1, (I.3.6.20)].

(7.3) Let S be a scheme, and G a reductive group scheme over S , and X a Noetherian S -scheme on which G acts trivially. Let M be a quasi-coherent (G, \mathcal{O}_X) -module. For (G, \mathcal{O}_X) -modules, see [Has4, Chapter 29]. Almost by definition, a (G, \mathcal{O}_X) -module and a $(G \times_S X, \mathcal{O}_X)$ -module (note that $G \times_S X$ is an X -group scheme) are the same thing.

We say that M is good if there is a Noetherian commutative ring R and a faithfully flat morphism of finite type $f : \mathrm{Spec} R \rightarrow X$ such that $G_R :=$

$\text{Spec } R \times_S G$ is a split reductive group scheme over R , and $\Gamma(\text{Spec } R, f^*M)$ is a good G_R -module. This notion is independent of the choice of f such that G_R is split reductive. When $X = \text{Spec } B$ is affine, then we also say that $\Gamma(X, M)$ is good, if M is good. If $g : X' \rightarrow X$ is a flat morphism of Noetherian schemes and M is a good quasi-coherent (G, \mathcal{O}_X) -module, then g^*M is good. If M is a quasi-coherent (G, \mathcal{O}_X) -module, g is faithfully flat, and g^*M is good, then M is good.

For a quasi-coherent (G, \mathcal{O}_X) -module M , we define the good locus of M to be

$$\text{Good}(M) = \{x \in X \mid M_x \text{ is a good } (\text{Spec } \mathcal{O}_{X,x} \times_S G)\text{-module}\}.$$

If $g : X' \rightarrow X$ is a flat morphism of Noetherian schemes, then $g^{-1}(\text{Good}(M)) = \text{Good}(g^*M)$. If $X = \text{Spec } R$ is affine, then for a (G, R) -module N , $\text{Good}(N)$ stands for $\text{Good}(\tilde{N})$, where \tilde{N} is the sheaf associated with N .

(7.4) Let the notation be as in (7.1).

For a poset ideal π of X^+ and a G -module M , we say that M belongs to π if $M_\lambda = 0$ for $\lambda \in X^+ \setminus \pi$.

Proposition 7.5. *Let π be a poset ideal of X^+ and M a G -module. Then the following are equivalent.*

- (1) M belongs to π .
- (2) For any R -finite subquotient N of M and any R -algebra K that is a field, $K \otimes_R N$ belongs to π .
- (3) For any R -finite subquotient N of M , any R -algebra K that is a field, and $\lambda \in X^+ \setminus \pi$, $\text{Hom}_G(\Delta_G(\lambda), K \otimes_R N) = 0$.
- (4) For any $\lambda \in X^+ \setminus \pi$, $\text{Hom}_G(\Delta_G(\lambda), M) = 0$.
- (5) M is a C_π -comodule, where C_π is the Donkin subcoalgebra of C with respect to π , see [Has1, (III.2.3.13)].

Proof. (1) \Rightarrow (2) is obvious.

(2) \Rightarrow (3) We may assume that $R = K$ and $N = M$. Then

$$\begin{aligned} \text{Hom}_G(\Delta_G(\lambda), M) &\cong \text{Hom}_G(M^*, \text{ind}_B^G(-w_0\lambda)) \\ &\cong \text{Hom}_B(M^*, -w_0\lambda) \cong \text{Hom}_B(w_0\lambda, M) \subset M_{w_0\lambda} = 0. \end{aligned}$$

(3) \Rightarrow (4) As M is the inductive limit of R -finite G -submodules of M , we may assume that M is R -finite. We use the Noetherian induction, and we may assume that the implication is true for R/I for any nonzero ideal I of R . If R is not a domain, then there is a nonzero ideal I of R such that the annihilator $0 : I$ of R is also nonzero. As $\text{Hom}_G(\Delta_G(\lambda), M/IM) = 0$ and $\text{Hom}_G(\Delta_G(\lambda), IM) = 0$, we have that $\text{Hom}_G(\Delta(\lambda), M) = 0$. So we may assume that R is a domain. Let N be the torsion part of M . Note that

$$0 \rightarrow N \rightarrow M \rightarrow K \otimes_R M$$

is exact, where K is the field of fractions of R . Hence N is a G -submodule of M . The annihilator of N is nontrivial, and hence $\text{Hom}_G(\Delta_G(\lambda), N) = 0$. On the other hand, by assumption, $\text{Hom}_G(\Delta_G(\lambda), K \otimes_R M) = 0$. So $\text{Hom}_G(\Delta_G(\lambda), M) = 0$, and we are done.

(4) \Rightarrow (5) is [Has1, (III.2.3.5)].

(5) \Rightarrow (1) As the coaction $\omega_M : M \rightarrow M' \otimes_R C_\pi$ is injective, it suffices to show that $M' \otimes_R C_\pi$ belongs to π , where M' is the R -module M with the trivial G -action. For this, it suffices to show that C_π belongs to π . This is proved easily by induction on the number of elements of π , if π is finite, almost by the definition of the Donkin system [Has1, (III.2.2)], and the fact that $\nabla_G(\lambda)$ belongs to π . Then the general case follows easily from the definition of C_π , see [Has1, (III.2.3.13)]. \square

Corollary 7.6. *Let M be a G -module, and π a poset ideal of the set of dominant weights X^+ . If M belongs to π , then $\text{Ext}_G^i(\Delta_G(\lambda), M) = 0$ for $i \geq 0$ and $\lambda \in X^+ \setminus \pi$.*

Proof. We use the induction on i . The case $i = 0$ is already proved in Proposition 7.5.

Let $i > 0$. Let C_π denote the Donkin subcoalgebra of $k[G]$. Consider the exact sequence

$$0 \rightarrow M \xrightarrow{\omega_M} M' \otimes_R C_\pi \rightarrow N \rightarrow 0.$$

Then N belongs to π , and $\text{Ext}_G^{i-1}(\Delta_G(\lambda), N) = 0$ by induction assumption. On the other hand, as C_π is good and R -finite R -projective by construction, $M' \otimes_R C_\pi$ is also good by [Has1, (III.4.1.8)]. Hence $\text{Ext}_G^i(\Delta_G(\lambda), M' \otimes_R C_\pi) = 0$. By the long exact sequence of the Ext-modules, we have that $\text{Ext}_G^i(\Delta_G(\lambda), M) = 0$. \square

Lemma 7.7. *Let the notation be as in (7.3). Let M be a coherent (G, \mathcal{O}_X) -module. Then $\text{Good}(M)$ is Zariski open in X .*

Proof. Let $f : \text{Spec } R \rightarrow X$ be a faithfully flat morphism of finite type such that G_R is split reductive. Let $M_R := \Gamma(\text{Spec } R, f^*M)$. Then $\text{Good}(M_R) = f^{-1}(\text{Good}(M))$. As f is a surjective open map, it suffices to show that $\text{Good}(M_R)$ is open in $\text{Spec } R$. So we may assume that $S = X = \text{Spec } R$ is affine and G is split, and we are to prove that $\text{Good}(N)$ is open for an R -finite G -module N .

As N is R -finite, there exists some finite poset ideal π of X^+ to which N belongs. Then $\text{Ext}_G^i(\Delta_G(\lambda), N) = 0$ for $\lambda \in X^+ \setminus \pi$ and $i \geq 0$. Set $L := \bigoplus_{\lambda \in \pi} \Delta_G(\lambda)$. Then $\text{Good}(N)$ is nothing but the complement of the support of the R -module $\text{Ext}_G^1(L, N)$ by [Has1, (III.2.3.8)]. As $\text{Ext}_G^1(L, N)$ is R -finite by [Has1, (III.2.3.19)], the support of $\text{Ext}_G^1(L, N)$ is closed, and we are done. \square

(7.8) Let the notation be as in (7.3). For a quasi-coherent (G, \mathcal{O}_X) -module M , the good dimension $\text{GD}(M)$ is defined to be $-\infty$ if $M = 0$. If $M \neq 0$ and there is an exact sequence

$$(5) \quad 0 \rightarrow M \rightarrow N_0 \rightarrow \cdots \rightarrow N_s \rightarrow 0$$

such that each N_i is good, then $\text{GD}(M)$ is defined to be the smallest s such that such an exact sequence exists. If there is no such an exact sequence, $\text{GD}(M)$ is defined to be ∞ .

(7.9) Assume that $X = \text{Spec } R$ is affine and G is split reductive. For a G -module M ,

$$\text{GD}(M) = \sup\{i \mid \bigoplus_{\lambda \in X^+} \text{Ext}_G^i(\Delta_G(\lambda), M) \neq 0\}.$$

Note that M is good if and only if $\text{GD}(M) \leq 0$. If $r \geq 0$, $s \geq -1$, and

$$0 \rightarrow M \rightarrow M_s \rightarrow \cdots \rightarrow M_0 \rightarrow N \rightarrow 0$$

is an exact sequence of G -modules with $\text{GD}(M_i) \leq i + r$, then $\text{GD}(M) \leq s + r + 1$ if and only if $\text{GD}(N) \leq r$.

If M and N are good and M is R -finite R -projective, then $M \otimes N$ is good, see [Has1, (III.4.5.10)]. Moreover, if M is R -finite R -projective with $\text{GD}(M) \leq s$, then M has an exact sequence of the form (5) such that each N_i is R -finite R -projective and good. Indeed, M belongs to some finite poset ideal π of X^+ , and when we truncate the cobar resolution of M as a C_π -comodule, then we obtain such a sequence.

It follows that for an R -finite R -projective G -module M , $\text{GD}(M) \leq s$ if and only if $\text{GD}(\kappa(\mathfrak{m}) \otimes_R M) \leq s$ for any maximal ideal \mathfrak{m} of R by [Has1, (III.4.1.8)].

It also follows that if $\text{GD}(M) \leq s$ and $\text{GD}(N) \leq t$ with M being R -finite R -projective, then $\text{GD}(M \otimes N) \leq s + t$.

Lemma 7.10. *Let V be an R -finite R -projective G -module with $\text{rank } V \leq n < \infty$. Then the following are equivalent.*

- (1) $\text{Sym } V$ is good.
- (2) $\bigoplus_{i=1}^{n-1} \text{Sym}_i V$ is good.
- (3) For $i = 1, \dots, n-1$, $\text{GD}(\bigwedge^i V) \leq i-1$.
- (4) For $i \geq 1$, $\text{GD}(\bigwedge^i V) \leq i-1$.

Proof. We may assume that R is a field.

(1) \Rightarrow (2) is trivial.

(2) \Rightarrow (3) We use the induction on i .

By assumption and the induction assumption, $\text{GD}(\text{Sym}_{i-j} V \otimes \bigwedge^j V) \leq j-1$ for $j = 1, \dots, i-1$. On the other hand, $\text{Sym}_i V$ is good. So by the exact sequence

$$(6) \quad 0 \rightarrow \bigwedge^i V \rightarrow \text{Sym}_1 V \otimes \bigwedge^{i-1} V \rightarrow \dots \rightarrow \text{Sym}_{i-1} V \otimes \bigwedge^1 V \rightarrow \text{Sym}_i V \rightarrow 0,$$

$\text{GD}(\bigwedge^i V) \leq i-1$.

(3) \Rightarrow (4) is trivial, as $\dim \bigwedge^i V \leq 1$ for $i \geq n$.

(4) \Rightarrow (1) Note that $\text{Sym}_0 V = R$ is good. Now use induction on $i \geq 1$ to prove that $\text{Sym}_i V$ is good (use the exact sequence (6) again). \square

Theorem 7.11. *Let S be a scheme, G a reductive S -group acting trivially on a Noetherian S -scheme X . Let M be a locally free coherent (G, \mathcal{O}_X) -module. Then*

$$(7) \quad \text{Good}(\text{Sym } M) = \{x \in X \mid \text{Sym}(\kappa(x) \otimes_{\mathcal{O}_{X,x}} M_x) \text{ is a good } (\text{Spec } \kappa(x) \times_S G)\text{-module}\},$$

and $\text{Good}(\text{Sym } M)$ is Zariski open in X .

Proof. Take a faithfully flat morphism of finite type $f : \text{Spec } R \rightarrow X$ such that $\text{Spec } R \times_S G$ is split reductive. Note that f is a surjective open map, and $f^{-1}(\text{Good}(\text{Sym } M)) = \text{Good}(\text{Sym } f^*M)$.

First we prove that $\text{Good}(\text{Sym } M)$ is open. We may assume that $S = X = \text{Spec } R$ is affine, and G is split reductive.

Then by Lemma 7.10 and Lemma 7.7,

$$\text{Good}(\text{Sym } M) = \text{Good}\left(\bigoplus_{i=1}^n \text{Sym}_i M\right)$$

is open, where the rank of M is less than or equal to n .

Next we prove that the equality (7) holds. Let $P \in \text{Spec } R$, and $x = f(P)$. Then $\text{Sym}(\kappa(x) \otimes_{\mathcal{O}_{X,x}} M_x)$ is good if and only if $\text{Sym}(\kappa(P) \otimes_{R_P} \Gamma(\text{Spec } R, f^*M)_P)$ is good. So we may assume that $S = X = \text{Spec } R$ is affine, and G is split reductive. Let N be an R -finite R -projective G -module of rank at most n . Then $(\text{Sym } N)_P$ is good if and only if $(\bigoplus_{i=1}^n \text{Sym}_i N)_P$ is good by Lemma 7.10. By [Has1, (III.4.1.8)], $(\bigoplus_{i=1}^n \text{Sym}_i N)_P$ is good if and only if $\kappa(P) \otimes_{R_P} (\bigoplus_{i=1}^n \text{Sym}_i N)_P$ is good. By Lemma 7.10 again, it is good if and only if $\kappa(P) \otimes_{R_P} (\text{Sym } N)_P$ is so. Thus the equality (7) was proved. \square

Corollary 7.12. *Let R be a Noetherian domain of characteristic zero, and G a reductive group over R . If M is an R -finite R -projective G -module, then $\{P \in \text{Spec } R \mid \text{Sym}(\kappa(P) \otimes_R M) \text{ is good}\}$ is a dense open subset of $\text{Spec } R$.*

Proof. By Theorem 7.11, it suffices to show that $\text{Good}(\text{Sym } M)$ is non-empty. But the generic point η of $\text{Spec } R$ is in $\text{Good}(\text{Sym } M)$. Indeed, $\kappa(\eta)$ is a field of characteristic zero, and any $\kappa(\eta) \otimes_R G$ -module is good. \square

REFERENCES

- [ABW] K. Akin, D. A. Buchsbaum, and J. Weyman, Schur functors and Schur complexes, *Adv. Math.* **44** (1982), 207–278.
- [AM] M. F. Atiyah and I. G. MacDonal, *Introduction to Commutative Algebra*, Addison-Wesley (1969).
- [Bt] J.-F. Boutot, Singularités rationnelles et quotients par les groupes réductifs, *Invent. Math.* **88** (1987), 65–68.

- [BH] W. Bruns and J. Herzog, *Cohen–Macaulay Rings*, first paper back ed., Cambridge (1998).
- [Cox] D. A. Cox, The homogeneous coordinate ring of a toric variety, *J. Algebraic Geom.* **4** (1995), 17–50.
- [DP] C. De Concini and C. Procesi, A characteristic free approach to invariant theory, *Adv. Math.* **21** (1976), 330–354.
- [Dol] I. Dolgachev, *Lectures on Invariant Theory*, *London Math. Soc. Lect. Note Series* **296**, Cambridge (2003).
- [Don1] S. Donkin, A filtration for rational modules, *Math. Z.* **177** (1981), 1–8.
- [Don2] S. Donkin, *Rational Representations of Algebraic Groups*, Lecture Notes in Math. **1140**, Springer (1985).
- [Don3] S. Donkin, Invariants of unipotent radicals, *Math. Z.* **198** (1988), 117–125.
- [Don4] S. Donkin, On tilting modules and invariants for algebraic groups, *Finite Dimensional Algebras and Related Topics* (Ottawa, ON, 1992), Kluwer (1994), pp. 59–77.
- [EKW] E. J. Elizondo, K. Kurano, and K.-i. Watanabe, The total coordinate ring of a normal projective variety, *J. Algebra* **276** (2004), 625–637.
- [Fr] E. M. Friedlander, A canonical filtration for certain rational modules, *Math. Z.* **188** (1985), 433–438.
- [GHKN] S. Goto, F. Hayasaka, K. Kurano, and Y. Nakamura, Rees algebras of the second syzygy module of the residue field of a regular local ring, *Commutative Algebra and Algebraic Geometry, Contemp. Math.* **390**, AMS (2005), pp. 97–108.
- [Grs1] F. D. Grosshans, The invariants of unipotent radicals of parabolic subgroups, *Invent. Math.* **73** (1983), 1–9.

- [Grs2] F. D. Grosshans, Contractions of the actions of reductive algebraic groups in arbitrary characteristic, *Invent. Math.* **107** (1992), 127–133.
- [Har] N. Hara, A characterization of rational singularities in terms of injectivity of Frobenius maps, *Amer. J. Math.* **120** (1998), 981–996.
- [Has1] M. Hashimoto, *Auslander–Buchweitz Approximations of Equivariant Modules*, *London Math. Soc. Lect. Note Series* **282**, Cambridge (2000).
- [Has2] M. Hashimoto, Good filtrations of symmetric algebras and strong F -regularity of invariant subrings, *Math. Z.* **236** (2001), 605–623.
- [Has3] M. Hashimoto, Surjectivity of multiplication and F -regularity of multigraded rings, *Commutative Algebra* (Grenoble/Lyon, 2001), *Contemp. Math.* **331** (2003), AMS, 153–170.
- [Has4] M. Hashimoto, Equivariant twisted inverses, *Foundations of Grothendieck Duality for Diagrams of Schemes* (J. Lipman, M. Hashimoto), *Lecture Notes in Math.* **1960**, Springer (2009), pp. 261–478.
- [Has5] M. Hashimoto, F -pure homomorphisms, strong F -regularity, and F -injectivity, *Comm. Algebra*, to appear.
- [Has6] M. Hashimoto, Good filtrations and F -purity of invariant subrings, *J. Math. Soc. Japan*, to appear.
- [Has7] M. Hashimoto, Equivariant total ring of fractions and factoriality of rings generated by semiinvariants, [arXiv:1009.5152](https://arxiv.org/abs/1009.5152)
- [HM] M. Hashimoto and M. Miyazaki, G -prime and G -primary G -ideals on G -schemes, [arXiv:0906.1441](https://arxiv.org/abs/0906.1441)
- [Hoc] M. Hochster, Foundations of tight closure theory, available at <http://www.math.lsa.umich.edu/~hochster/mse.html>
- [HH1] M. Hochster and C. Huneke, Tight closure and strong F -regularity, *Mém. Soc. Math. France* **38** (1989), 119–133.

- [HH2] M. Hochster and C. Huneke, Tight closure, invariant theory, and the Briançon–Skoda theorem, *J. Amer. Math. Soc.* **3** (1990), 31–116.
- [HH3] M. Hochster and C. Huneke, F -regularity, test elements, and smooth base change, *Trans. Amer. Math. Soc.* **346** (1994), 1–62.
- [HH4] M. Hochster and C. Huneke, Tight closure of parameter ideals and splitting in module-finite extensions, *J. Algebraic Geom.* **3** (1994), 599–670.
- [HR] M. Hochster and J. Roberts, Rings of invariants of reductive groups acting on regular rings are Cohen–Macaulay, *Adv. Math.* **13** (1974), 115–175.
- [Hum] J. E. Humphreys, *Linear Algebraic Groups*, third printing, *Graduate Texts in Math.* **21** (1987).
- [Jan] J. C. Jantzen, *Representations of Algebraic Groups*, 2nd ed., AMS (2003).
- [Kem] G. Kemper, A characterization of linearly reductive groups by their invariants, *Transform Groups*, **5** (2000), 1–8.
- [KKMS] G. Kempf, F. Knudsen, D. Mumford and B. Saint-Donat, *Toroidal Embeddings I*, *Lecture Notes in Math.* **339**, Springer (1973).
- [Kun] E. Kunz, On Noetherian rings of characteristic p , *Amer. J. Math.* **98** (1976), 999–1013.
- [LS] G. Lyubeznik and K. E. Smith, Strong and weak F -regularity are equivalent for graded rings, *Amer. J. Math.* **121** (1999), 1279–1290.
- [Mat] O. Mathieu, Filtrations of G -modules, *Ann. Sci. École Norm. Sup.* (4) **23** (1990), 625–644.
- [Miy] M. Miyazaki, Invariants of the unipotent radical of a Borel subgroup, *The 29th Symposium on Commutative Algebra in Japan* (Nagoya, 2007), pp. 43–50, <http://www.math.nagoya-u.ac.jp/~hasimoto/commalg-proc.html>

- [MFK] D. Mumford, J. Fogarty and F. Kirwan, *Geometric Invariant Theory*, third edition, Springer (1994).
- [Mur] M. P. Murthy, A note on factorial rings, *Arch. Math.* **15** (1964), 418–420.
- [Pop1] V. L. Popov, On the stability of the action of an algebraic group on an algebraic variety, *Math. USSR-Izv.* **6** (1972), 367–379.
- [Pop2] V. L. Popov, The Picard group of homogeneous spaces of linear algebraic groups and one-dimensional homogeneous vector bundles, *Math. USSR-Izv.* **8** (1975), 301–327.
- [Pop3] V. L. Popov, Contraction of the actions of reductive algebraic groups, *Math. USSR-Sb.* **58** (1987), 311–335.
- [RR] S. Ramanan and A. Ramanathan, Projective normality of flag varieties and Schubert varieties, *Invent. Math.* **79** (1985), 217–224.
- [Ros] M. Rosenlicht, Toroidal algebraic groups, *Proc. Amer. Math. Soc.* **12** (1961), 984–988.
- [SvdB] K. E. Smith and M. Van den Bergh, Simplicity of rings of differential operators in prime characteristic, *Proc. London Math. Soc.* **75** (1997), 32–62.
- [Vel] J. D. Vélez, Openness of the F -rational locus and smooth base change, *J. Algebra* **172** (1995), 425–453.